

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Refined semiclassical asymptotics for fractional
powers of the Laplace operator

Rupert L. Frank, Leander Geisinger

Preprint 2011/020

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Refined semiclassical asymptotics for fractional
powers of the Laplace operator

Rupert L. Frank, Leander Geisinger

Preprint 2011/020

Fachbereich Mathematik
Fakultät Mathematik und Physik
Universität Stuttgart
Pfaffenwaldring 57
D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de

WWW: <http://www.mathematik.uni-stuttgart.de/preprints>

ISSN **1613-8309**

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.

LaTeX-Style: Winfried Geis, Thomas Merkle

1. INTRODUCTION AND MAIN RESULT

1.1. **Introduction.** In this paper we study the asymptotic behavior of eigenvalues for fractional powers of the Laplacian. The operator $(-\Delta)^s$ with $0 < s < 1$ appears in numerous fields of mathematical physics, mathematical biology and mathematical finance and has attracted a lot of attention recently. The key difference between this operator and the usual Laplacian is the non-locality of $(-\Delta)^s$, which allows one to model long-range interactions in applications and leads to challenging mathematical problems.

From a probabilistic point of view, the fractional Laplacian of order s on a domain $\Omega \subset \mathbb{R}^d$ can be defined as the generator of the $2s$ -stable process killed upon exiting Ω . A more operator theoretic definition, which we employ here, is in terms of the quadratic form

$$C_{s,d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|u(x) - u(y)|^2}{|x - y|^{d+2s}} dx dy = \int_{\mathbb{R}^d} |p|^{2s} |\hat{u}(p)|^2 dp, \quad (1)$$

restricted to functions $u \in H^s(\mathbb{R}^d)$ which satisfy $u \equiv 0$ in $\mathbb{R}^d \setminus \bar{\Omega}$. Here $H^s(\mathbb{R}^d)$ is the Sobolev space of order s , $\hat{u}(p) = (2\pi)^{-d/2} \int e^{-ip \cdot x} u(x) dx$ is the Fourier transform of u and $C_{s,d}$ is an explicit constant given in (5). The identity in (1) is an easy consequence of Plancherel's theorem.

For bounded domains Ω the spectrum of the fractional Laplacian is discrete and we denote its eigenvalues (in increasing order, repeated according to multiplicities) by $\lambda_n^{(s)}$. Our main result in this paper is a two-term asymptotic expansion of the sum of these eigenvalues,

$$\frac{1}{N} \sum_{n=1}^N \lambda_n^{(s)} = C_{d,s}^{(1)} |\Omega|^{-2s/d} N^{2s/d} + C_{d,s}^{(2)} |\partial\Omega| |\Omega|^{-(d-1+2s)/d} N^{(2s-1)/d} (1 + o(1)) \quad (2)$$

as $N \rightarrow \infty$. Here $|\Omega|$ and $|\partial\Omega|$ denote the d -dimensional measure of Ω and the $(d-1)$ -dimensional surface measure of $\partial\Omega$, respectively, and $C_{d,s}^{(1)}$ and $C_{d,s}^{(2)}$ are positive, universal constants, depending only on d and s , for which we shall obtain explicit expressions. Our result is valid for non-smooth domains, requiring only that $\partial\Omega \in C^{1,\alpha}$ for some (arbitrarily small) $\alpha > 0$. It is remarkable that, despite the fact that we are dealing with a non-local operator, both coefficients in (2) have a local form, depending only on Ω and $\partial\Omega$, just like in the case of the Laplacian. This will become clearer from the reformulation given in Theorem 1 below.

In order to avoid confusion, we emphasize that the fractional Laplacian of order s on a domain Ω is different from the Dirichlet Laplacian on Ω raised to the s -th power. For the Dirichlet Laplacian, and hence for its fractional powers, asymptotics analogous to (2) are well-known. One of our results is that, while the first terms in (2) coincide for both operators, the second terms do *not*. This means, in particular, that our result cannot be obtained from the study of the (local) Dirichlet Laplacian, and that our analysis needs to take into account the non-locality inherent in (2). For further results about the relation between the fractional Laplacian on a domain and the fractional power of the Dirichlet Laplacian we refer to [CS05]; see also Section 6 below.

The one-term asymptotics $\lambda_N^{(s)} = \frac{d+2s}{d} C_{d,s}^{(1)} |\Omega|^{-2s/d} N^{2s/d} (1 + o(1))$, which is a fractional version of Weyl's law, is a classical result of Blumenthal and Gettoor [BG59]. More recently, Bañuelos and Kulczycki [BK08] and Bañuelos, Kulczycki and Siudeja [BKS09] have shown a two-term asymptotic formula for $\sum_{n=1}^{\infty} \exp(-t\lambda_n^{(s)})$ as $t \rightarrow 0$. Note that $\sum_{n=1}^{\infty} \exp(-t\lambda_n^{(s)})$ and $\frac{1}{N} \sum_{n=1}^N \lambda_n^{(s)}$ correspond to the Abel and Cesàro summation of the sequence $\lambda_n^{(s)}$, respectively. As is well-known, asymptotics of Cesàro means imply asymptotics of Abel means, but not vice versa. Hence for $C^{1,\alpha}$ domains we recover and improve upon the result of [BK08, BKS09].

This is, actually, a significant improvement since our asymptotics are no longer derived for the infinitely smooth function e^{-tE} of the fractional Laplacian, but, as we shall see shortly, for the Lipschitz function $(\Lambda - E)_+$. Moreover, since we are no longer able to apply the probabilistic machinery available for the partition function, we have to find new and more robust tools. Our methods also work for the ordinary Dirichlet Laplacian on a bounded domain, and in [FG11] we use the techniques developed here to give an elementary and short proof of two-term asymptotics in that case.

Another point in which we go beyond [BK08, BKS09] is that we give an expression for the constant $C_{d,s}^{(2)}$ in (2) in terms of a model operator on a *half-line* instead of a model operator on a *half-space*. In this way our expression is similar to familiar two-term formulas in semiclassical analysis; see, for instance, [SV97]. This is possible due to some recent beautiful results of Kwaśnicki [Kwa10b] about a general class of half-line operators.

We find it convenient to prove (2) in an equivalent form, namely

$$\sum_{n=1}^{\infty} (\Lambda - \lambda_n^{(s)})_+ = L_{s,d}^{(1)} |\Omega| \Lambda^{1+d/2s} - L_{s,d}^{(2)} |\partial\Omega| \Lambda^{1+(d-1)/2s} (1 + o(1)) \quad \text{as } \Lambda \rightarrow \infty. \quad (3)$$

Here $x_+ := \max\{x, 0\}$ denotes the positive part of a number x . (The fact that (2) and (3) are equivalent is well-known to experts in the field, but we include a short proof in Section 7 for the sake of completeness, see Lemma 20.) Note also that (3) can be rewritten as

$$\sum_{n=1}^{\infty} (1 - h^{2s} \lambda_n^{(s)})_+ = L_{s,d}^{(1)} |\Omega| h^{-d} - L_{s,d}^{(2)} |\partial\Omega| h^{-d+1} (1 + o(1)) \quad \text{as } h \rightarrow 0+, \quad (4)$$

and this is the form in which we shall state and prove our main theorem. The small parameter h has the interpretation of Planck's constant and (4) emphasizes the semiclassical nature of the problem.

Our approach extends the multiscale analysis to the fractional setting. By this we mean that we localize simultaneously on different length scales according to the distance from the boundary. Of course, a main difficulty when dealing with our non-local operator comes from the treatment of the localization error. At this point we have to improve upon previous results from [LY88, SSS10]. Another major impass, as compared to the local case, is the analysis of a one-dimensional model operator for which an (almost) explicit diagonalization is far from trivial. This is where Kwaśnicki's work [Kwa10b] enters. It requires, however, still substantial work to bring these results into a form which is useful for us. We will

explain the strategy of our proof in more detail in Subsection 1.3 after a precise statement of our main result.

Throughout this paper we assume that the dimension $d \geq 2$. In the one dimensional case (the fractional Laplacian on an interval) considerably stronger results are known [KKMS10, Kwa10a]. The powerful methods developed there are, however, intrinsically one-dimensional and seem of little help in the multi-dimensional case. The question raised in [BKS09] of whether an analogue of Ivrii's two-term asymptotics [Ivr80] holds for $\lambda_n^{(s)}$ in $d \geq 2$ without Abel or Cesàro averaging remains a challenging open problem.

1.2. Main Result. Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$, be a bounded open set. For $h > 0$ and $0 < s < 1$ let

$$H_\Omega = (-h^2 \Delta)^s - 1$$

be the self-adjoint operator in $L^2(\Omega)$ generated by the quadratic form

$$(u, H_\Omega u) = \int_{\mathbb{R}^d} (|hp|^{2s} - 1) |\hat{u}(p)|^2 dp$$

with form domain

$$\mathcal{H}^s(\Omega) = \{u \in H^s(\mathbb{R}^d) : u \equiv 0 \text{ on } \mathbb{R}^d \setminus \bar{\Omega}\}.$$

For $0 < s < 1$ we have the representation

$$(u, H_\Omega u) = C_{s,d} h^{2s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|u(x) - u(y)|^2}{|x - y|^{d+2s}} dx dy - \int_{\Omega} |u(x)|^2 dx$$

with constant

$$C_{s,d} = 2^{2s-1} \pi^{-d/2} \frac{\Gamma(d/2 + s)}{|\Gamma(-s)|} > 0. \quad (5)$$

Our main results hold without any global geometric conditions on Ω . We only require weak smoothness conditions on the boundary - namely that the boundary belongs to the class $C^{1,\alpha}$ for some $\alpha > 0$. That is, the local charts of $\partial\Omega$ are differentiable and the derivatives are Hölder continuous with exponent α .

Theorem 1. *Let $0 < s < 1$ and assume that the boundary of Ω satisfies $\partial\Omega \in C^{1,\alpha}$ with some $0 < \alpha \leq 1$. Then*

$$\text{Tr}(H_\Omega)_- = L_{s,d}^{(1)} |\Omega| h^{-d} - L_{s,d}^{(2)} |\partial\Omega| h^{-d+1} + R_h \quad (6)$$

with $R_h = o(h^{-d+1})$ as $h \rightarrow 0+$. Here

$$L_{s,d}^{(1)} = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} (|p|^{2s} - 1)_- dp \quad (7)$$

and the positive constant $L_{s,d}^{(2)}$ is given in (28).

More precisely, we have the lower bound $R_h \geq -Ch^{-d+1+\epsilon_-}$ for any

$$0 < \epsilon_- < \begin{cases} \frac{\alpha}{\alpha+2} & \text{if } 1/2 \leq s < 1, \\ \frac{2s\alpha}{\alpha+1+2s} & \text{if } 0 < s < 1/2, \end{cases}$$

and the upper bound $R_h \leq Ch^{-d+1+\epsilon_+}$ for any

$$\begin{aligned} 0 < \epsilon_+ < \frac{\alpha}{\alpha + 2} & \quad \text{if } 1 - d/4 \leq s < 1, \\ 0 < \epsilon_+ \leq \frac{\alpha(2s - 1 + d/2)}{\alpha + 2s + d/2} & \quad \text{if } 0 < s < 1 - d/4. \end{aligned}$$

We do not claim that our remainder estimates are sharp. They show, however, that our methods are rather explicit and they correctly reflect the intuitive fact that the estimate worsens as the boundary gets rougher. We also mention that for not too small s we (almost) get the same remainder estimate $h^{-d+1+\alpha/(\alpha+2)}$ that our method yields in the local case $s = 1$ [FG11].

In Section 6 we will derive several representations of the constant $L_{s,d}^{(2)}$ in (6). One of these, which emphasizes the semi-classical nature of the problem, leads to a rewriting of (6) as

$$\mathrm{Tr}(H_\Omega)_- = \iint_{T\Omega} (|p|^{2s} - 1)_- \frac{dpdx}{(2\pi h)^d} - \iint_{T\partial\Omega} \zeta(|p'|^{-2s}) \frac{dp'd\sigma(x)}{(2\pi h)^{d-1}} + R_h, \quad (8)$$

where $T\Omega = \Omega \times \mathbb{R}^d$ and $T\partial\Omega = \partial\Omega \times \mathbb{R}^{d-1}$ are the cotangent bundles over Ω and $\partial\Omega$, respectively, and where $d\sigma$ is the surface element of $\partial\Omega$. Here ζ is a universal (i.e., depending on s , but independent of Ω or d) function, which has the interpretation of an energy shift (the integral of a spectral shift). It is given in terms of a one-dimensional model operator A^+ on the half-line \mathbb{R}_+ and its analogue A on the whole line (see Section 3) by

$$\zeta(\mu) = \mu^{-1} \int_0^\infty (a(t, t, \mu) - a^+(t, t, \mu)) dt, \quad \mu > 0,$$

where $a(t, u, \mu)$ and $a^+(t, u, \mu)$ denote the integral kernels of $(A - \mu)_-$ and $(A^+ - \mu)_-$, respectively. Another representation, derived in Remark 6.1, shows that our result is consistent with the result of [BK08, BKS09].

In Section 6 we also prove that

$$L_{s,d}^{(2)} > 0.$$

Moreover, we compare this constant with the one obtained from the corresponding fractional power of the Dirichlet Laplacian.

Proposition 2. *Let $0 < s < 1$ and assume that the boundary of Ω satisfies $\partial\Omega \in C^{1,\alpha}$ with some $0 < \alpha \leq 1$. Let $-\Delta_\Omega$ be the Dirichlet Laplacian on Ω . Then*

$$\mathrm{Tr} \left((-h^2 \Delta_\Omega)^s - 1 \right)_- = L_{s,d}^{(1)} |\Omega| h^{-d} - \tilde{L}_{s,d}^{(2)} |\partial\Omega| h^{-d+1} + R_h \quad (9)$$

with $R_h = o(h^{-d+1})$ as $h \rightarrow 0+$. Here $L_{s,d}^{(1)}$ is the same as in (7) and $\tilde{L}_{s,d}^{(2)}$ satisfies

$$L_{s,d}^{(2)} < \tilde{L}_{s,d}^{(2)}. \quad (10)$$

In other words, the operators H_Ω and $(-h^2 \Delta_\Omega)^s - 1$ differ semi-classically to first sub-leading order.

1.3. Strategy of the proof. The proof of Theorem 1 is divided into three main steps: First, we localize the operator H_Ω into balls, whose size varies depending on the distance to the complement of Ω . Then we can analyze separately the semiclassical limit in the bulk and at the boundary.

The key idea is to choose the localization depending on the distance to the complement of Ω , see [Hör85, Theorem 17.1.3] and [SS03]. Let $d(u) = \inf\{|x - u| : x \notin \Omega\}$ denote the distance of $u \in \mathbb{R}^d$ to the complement of Ω . We set

$$l(u) = \frac{1}{2} \left(1 + (d(u)^2 + l_0^2)^{-1/2} \right)^{-1}, \quad (11)$$

where $0 < l_0 \leq 1/2$ is a small parameter depending only on h . Indeed, we will finally choose l_0 proportional to h^β with suitable $0 < \beta < 1$.

In Section 5 we construct real-valued functions $\phi_u \in C_0^\infty(\mathbb{R}^d)$ with support in the ball $B_u = \{x \in \mathbb{R}^d : |x - u| < l(u)\}$. For all $u \in \mathbb{R}^d$ these functions satisfy

$$\|\phi_u\|_\infty \leq C, \quad \|\nabla \phi_u\|_\infty \leq C l(u)^{-1} \quad (12)$$

and for all $x \in \mathbb{R}^d$

$$\int_{\mathbb{R}^d} \phi_u^2(x) l(u)^{-d} du = 1. \quad (13)$$

Here and in the following the letter C denotes various positive constants that are independent of u , l_0 and h .

Proposition 3. *There is a constant $C > 0$ depending only on s and d such that for all $0 < l_0 \leq 1/2$ and all $0 < h \leq C^{-1}l_0$ the estimates*

$$0 \leq \text{Tr}(H_\Omega)_- - \int_{\mathbb{R}^d} \text{Tr}(\phi_u H_\Omega \phi_u)_- l(u)^{-d} du \leq C h^{-d+2} l_0^{-1} R_{\text{loc}}(h, l_0)$$

hold with a remainder

$$R_{\text{loc}}(h, l_0) = \begin{cases} 1 & \text{if } 1 - d/4 < s < 1 \\ |\ln(l_0/h)|^{1/2} & \text{if } 0 < s = 1 - d/4 \\ (l_0/h)^{2-2s-d/2} & \text{if } 0 < s < 1 - d/4 \end{cases}.$$

In view of this result, one can analyze the local asymptotics, i.e., the asymptotic behavior of $\text{Tr}(\phi_u H_\Omega \phi_u)_-$, separately on different parts of Ω . First, we consider the bulk, where the influence of the boundary is not felt.

Proposition 4. *Assume that $\phi \in C_0^1(\Omega)$ is supported in a ball of radius $l > 0$ and that*

$$\|\nabla \phi\|_\infty \leq C l^{-1}. \quad (14)$$

Then for all $h > 0$ the estimates

$$-C l^{d-2} h^{-d+2} \leq \text{Tr}(\phi H_\Omega \phi)_- - L_{s,d}^{(1)} \int_\Omega \phi^2(x) dx h^{-d} \leq 0$$

hold with a constant depending only on the constant in (14).

Close to the boundary of Ω , more precisely, if the support of ϕ intersects the boundary, a boundary term of the order h^{-d+1} appears.

Proposition 5. *Assume that $\phi \in C_0^1(\mathbb{R}^d)$ is supported in a ball of radius $0 < l \leq 1$ intersecting the boundary of Ω and assume that (14) is satisfied. Then for all $h > 0$ the estimates*

$$-\tilde{R}_{\text{bd}}(l, h) \leq \text{Tr}(\phi H_\Omega \phi)_- - L_{s,d}^{(1)} \int_\Omega \phi^2(x) dx h^{-d} + L_{s,d}^{(2)} \int_{\partial\Omega} \phi^2(x) d\sigma(x) h^{-d+1} \leq R_{\text{bd}}(l, h)$$

hold. Here $d\sigma$ denotes the $(d-1)$ -dimensional volume element of $\partial\Omega$ and the remainder terms satisfy for any $0 < \delta_1 < 1$ and $0 < \delta_2 < \min\{1, 2s\}$

$$\begin{aligned} R_{\text{bd}}(l, h) &\leq C_{\delta_1} \left(\frac{l^{d-1-\delta_1}}{h^{d-1-\delta_1}} + \frac{l^{d+\alpha}}{h^d} \right), \\ \tilde{R}_{\text{bd}}(l, h) &\leq C_{\delta_1, \delta_2} \left(\frac{l^{d-1-\delta_1}}{h^{d-1-\delta_1}} + \frac{l^{d-1-\delta_2}}{h^{d-1-\delta_2}} + \frac{l^{2\alpha+d-1}}{h^{d-1}} + \frac{l^{d+\alpha}}{h^d} \right), \end{aligned}$$

with constants depending on $\delta_1, \delta_2, \Omega, \|\phi\|_\infty$ and the constant in (14).

Based on these propositions we can complete the proof of Theorem 1.

Proof of Theorem 1. In order to apply Proposition 5 to the operators $\phi_u H_\Omega \phi_u$, we need to estimate $l(u)$ uniformly. Let

$$U(\Omega) = \{u \in \mathbb{R}^d : B_u \cap \partial\Omega \neq \emptyset\}$$

be a small neighborhood of the boundary. For $u \in U(\Omega)$ we have $d(u) \leq l(u)$, which by the definition of $l(u)$ implies

$$l(u) \leq l_0/\sqrt{3}. \quad (15)$$

In view of (12) and (15) we can apply Proposition 4 and Proposition 5 to all functions $\phi_u, u \in \mathbb{R}^d$, if l_0 is sufficiently small. Combining these results with Proposition 3 we get

$$\begin{aligned} &-C \int_{\Omega \setminus U(\Omega)} l(u)^{-2} du h^{-d+2} - \int_{U(\Omega)} \tilde{R}_{\text{bd}}(l(u), h) l(u)^{-d} du \\ &\leq \text{Tr}(H_\Omega)_- - L_{s,d}^{(1)} \int_{\mathbb{R}^d} \int_\Omega \phi_u^2(x) dx \frac{du}{l(u)^d} h^{-d} + L_{s,d}^{(2)} \int_{\mathbb{R}^d} \int_{\partial\Omega} \phi_u^2(x) d\sigma(x) \frac{du}{l(u)^d} h^{-d+1} \\ &\leq \int_{U(\Omega)} R_{\text{bd}}(l(u), h) l(u)^{-d} du + Ch^{-d+2} l_0^{-1} R_{\text{loc}}(l_0, h). \end{aligned}$$

Now we change the order of integration and in view of (13) we obtain

$$\begin{aligned} &-C \int_{\Omega \setminus U(\Omega)} l(u)^{-2} du h^{-d+2} - \int_{U(\Omega)} \tilde{R}_{\text{bd}}(l(u), h) l(u)^{-d} du \\ &\leq \text{Tr}(H_\Omega)_- - L_{s,d}^{(1)} |\Omega| h^{-d} + L_{s,d}^{(2)} |\partial\Omega| h^{-d+1} \\ &\leq \int_{U(\Omega)} R_{\text{bd}}(l(u), h) l(u)^{-d} du + Ch^{-d+2} l_0^{-1} R_{\text{loc}}(l_0, h). \end{aligned} \quad (16)$$

It remains to estimate the error terms.

By definition of $l(u)$ we have

$$l(u) \geq \frac{1}{4} \min(d(u), 1) \quad \text{and} \quad l(u) \geq \frac{l_0}{4} \quad (17)$$

for all $u \in \mathbb{R}^d$. For $u \in \Omega \setminus U(\Omega)$, we find $d(u) \geq l(u) \geq l_0/4$. Hence, we can estimate

$$\int_{\Omega \setminus U(\Omega)} l(u)^{-2} du \leq C \left(1 + \int_{\{d(u) \geq l_0/4\}} d(u)^{-2} du \right) \leq C \left(1 + \int_{l_0/4}^{\infty} t^{-2} |\partial\Omega_t| dt \right),$$

where $|\partial\Omega_t|$ denotes the surface area of the boundary of $\Omega_t = \{x \in \Omega : d(x) > t\}$. Using the fact that $|\partial\Omega_t|$ is uniformly bounded and that $|\partial\Omega_t| = 0$ for large t , we get

$$\int_{\Omega \setminus U(\Omega)} l(u)^{-2} du \leq Cl_0^{-1}. \quad (18)$$

For $u \in U(\Omega)$ the inequalities (15) and (17) show that $l(u)$ is proportional to l_0 . Since $B_u \cap \partial\Omega \neq \emptyset$ we find $d(u) < l(u) \leq Cl_0$ and

$$\int_{U(\Omega)} l(u)^a du \leq Cl_0^a \int_{\{d(u) \leq l_0\}} du \leq Cl_0^{a+1}, \quad (19)$$

for any $a \in \mathbb{R}$.

We insert (18) and (19) into (16) and get (using the fact that $h \leq C^{-1}l_0$)

$$\begin{aligned} -C (l_0^{-\delta_2} h^{\delta_2} + l_0^{2\alpha} + l_0^{\alpha+1} h^{-1}) &\leq h^{d-1} \left(\text{Tr}(H_\Omega)_- - L_{s,d}^{(1)} |\Omega| h^{-d} + L_{s,d}^{(2)} |\partial\Omega| h^{-d+1} \right) \\ &\leq C (l_0^{-\delta_1} h^{\delta_1} + l_0^{\alpha+1} h^{-1} + l_0^{-1} h R_{\text{loc}}(l_0, h)). \end{aligned} \quad (20)$$

In order to choose l_0 we need to distinguish several cases. For the lower bound we recall that $0 < \delta_2 < \min\{1, 2s\}$. The stated lower bound on R_h follows with l_0 proportional to h^β , where $\beta = (1 + \delta_2)/(1 + \alpha + \delta_2)$.

For the upper bound we have $0 < \delta_1 < 1$. If $1 - d/4 < s < 1$, we pick l_0 proportional to h^β , where $\beta = (1 + \delta_1)/(1 + \alpha + \delta_1)$. If $0 < s \leq 1 - d/4$, we pick h^β , where $\beta = (2s + d/2)/(\alpha + 2s + d/2)$. This completes the proof of Theorem 1. \square

The remainder of the text is structured as follows. First we analyze the local asymptotics in the bulk and prove Proposition 4. This is done in Section 2. In Section 3 we consider the local asymptotics in the case where Ω is replaced by a half-space. We reduce the problem close to the boundary to the analysis of a one-dimensional model operator given on a half-line and give an analogue of Proposition 5 for a half-space. In Section 4 we show how Proposition 5 follows from the previous considerations by local straightening of the boundary. In Section 5, we perform the localization and, in particular, prove Proposition 3. In Section 8 we provide some technical results about the one-dimensional model operator introduced in Section 3.

Notation. We define the positive and negative parts of a real number x by $x_\pm = \max\{0, \pm x\}$. We use a similar notation for the heavy side function, namely, $x_\pm^0 = 1$ if $\pm x \geq 0$ and $x_\pm^0 = 0$ if $\pm x < 0$. For a self-adjoint operator X , the operators X_\pm and X_\pm^0 are defined similarly via the spectral theorem.

2. LOCAL ASYMPTOTICS IN THE BULK

This section is a warm-up dealing with the spectral asymptotics in the boundaryless case. Although the estimates in this case are essentially known, we include a proof for the sake of completeness and in order to introduce the methods that will be important later on. We divide the proof of Proposition 4 into two subsections containing the lower and the upper bound, respectively. The operator

$$H_0 = (-h^2\Delta)^s - 1 \quad \text{in } L^2(\mathbb{R}^d),$$

defined with form domain $H^s(\mathbb{R}^d)$, will appear frequently.

2.1. Lower bound. The lower bound is given by a variant of the Berezin-Lieb-Li-Yau inequality, see [Ber72, Lie73, LY83]. For later purposes we record this as

Lemma 6. *For any $\phi \in L^2(\mathbb{R}^d)$ and $h > 0$*

$$\mathrm{Tr}(\phi H_\Omega \phi)_- \leq L_{s,d}^{(1)} \int_{\mathbb{R}^d} \phi^2(x) dx h^{-d}.$$

Proof. We apply the variational principle for the sum of the eigenvalues

$$-\mathrm{Tr}(\phi H_\Omega \phi)_- = \inf_{0 \leq \gamma \leq 1} \mathrm{Tr}(\gamma \phi H_\Omega \phi),$$

where the infimum is taken over all trial density matrices, i.e., over all trace-class operators $0 \leq \gamma \leq 1$ with range belonging to the form domain of H_Ω . We apply this twice and find

$$\mathrm{Tr}(\phi H_\Omega \phi)_- \leq \mathrm{Tr}(\phi H_0 \phi)_- \leq \mathrm{Tr}(\phi (H_0)_- \phi).$$

Applying the Fourier transform to diagonalize the operator $(H_0)_-$ yields the bound

$$\mathrm{Tr}(\phi (H_0)_- \phi) = \frac{1}{(2\pi h)^d} \iint \phi(x)^2 (|p|^{2s} - 1)_- dp dx = L_{s,d}^{(1)} \int \phi(x)^2 dx h^{-d},$$

as claimed. □

2.2. Upper bound. We now assume that ϕ satisfies the conditions of Proposition 4. In particular, we assume that ϕ has support in Ω . To derive the upper bound we put $\gamma = (H_0)_-$, i.e.,

$$\gamma(x, y) = (2\pi h)^{-d} \int_{|p| < 1} e^{ip \cdot (x-y)/h} dp,$$

and obtain that

$$\begin{aligned} -\mathrm{Tr}(\phi H_\Omega \phi)_- &\leq \mathrm{Tr}(\gamma \phi H_\Omega \phi) = \mathrm{Tr}(\gamma \phi H_0 \phi) \\ &= \int_{|p| < 1} (\|(-h^2\Delta)^{s/2} \phi e^{ip \cdot /h}\|_2^2 - \|\phi\|_2^2) \frac{dp}{(2\pi h)^d}. \end{aligned} \quad (21)$$

Lemma 7. *For $\phi \in C_0^\infty(\mathbb{R}^d)$ and $h > 0$ we have*

$$\|(-h^2\Delta)^{s/2} \phi e^{-ip \cdot /h}\|_2^2 = |p|^{2s} \|\phi\|_2^2 + \int \left(\frac{1}{2} (|p + h\eta|^{2s} + |p - h\eta|^{2s}) - |p|^{2s} \right) |\hat{\phi}(\eta)|^2 d\eta.$$

Proof. By Plancherel's theorem we get

$$\begin{aligned}
& \|(-h^2\Delta)^{s/2}\phi e^{ip\cdot/h}\|_2^2 \\
&= \iiint |\xi|^{2s} \phi(x)\phi(y) e^{i(p-\xi)\cdot(x-y)/h} \frac{dx dy d\xi}{(2\pi h)^d} \\
&= \frac{1}{2} \iiint |\xi|^{2s} (\phi^2(x) + \phi^2(y) - |\phi(x) - \phi(y)|^2) e^{i(p-\xi)\cdot(x-y)/h} \frac{dx dy d\xi}{(2\pi h)^d}. \tag{22}
\end{aligned}$$

In the first two terms we perform the ξ integration and either the x or the y integration to arrive at

$$\frac{1}{2} \iiint |\xi|^{2s} (\phi^2(x) + \phi^2(y)) e^{i(p-\xi)\cdot(x-y)/h} \frac{dx dy d\xi}{(2\pi h)^d} = |p|^{2s} \int \phi^2(x) dx. \tag{23}$$

We are left with calculating the third term in (22). Again, by Plancherel's theorem we see that it equals

$$\frac{1}{2} \iiint |\xi|^{2s} \left| \hat{\phi}\left(\frac{\eta}{h}\right) \right|^2 |1 - e^{-iz\cdot\eta/h}|^2 e^{i(p-\xi)\cdot z/h} \frac{d\eta dz d\xi}{(2\pi)^d h^{2d}}.$$

We can write

$$|1 - e^{-iz\cdot\eta/h}|^2 = 2 - e^{iz\cdot\eta/h} - e^{-iz\cdot\eta/h}$$

and perform the integration in z and ξ to obtain

$$\begin{aligned}
& \frac{1}{2} \iiint |\xi|^{2s} |\phi(x) - \phi(y)|^2 e^{i(p-\xi)\cdot(x-y)/h} \frac{dx dy d\xi}{(2\pi h)^d} \\
&= \frac{1}{h^d} \int \left(|p|^{2s} - \frac{1}{2} (|p + \eta|^{2s} + |p - \eta|^{2s}) \right) \left| \hat{\phi}\left(\frac{\eta}{h}\right) \right|^2 d\eta. \tag{24}
\end{aligned}$$

Hence, combining (22), (23) and (24) yields the claim. \square

In view of identity (21) and Lemma 7 we conclude

$$\text{Tr}(\gamma\phi H_0\phi) = (2\pi h)^{-d} \int_{|p|<1} (|p|^{2s} - 1) dp \|\phi\|_2^2 + (2\pi h)^{-d} \int_{|p|<1} R_h(p) dp \tag{25}$$

with

$$R_h(p) = \int \left(\frac{1}{2} (|p + h\eta|^{2s} + |p - h\eta|^{2s}) - |p|^{2s} \right) |\hat{\phi}(\eta)|^2 d\eta.$$

We proceed to estimate $R_h(p)$. Note that for any $a > 0$

$$\max_{|t|\leq a} ((a+t)^s + (a-t)^s) = 2a^s.$$

Taking $a = |p|^2 + |\eta|^2$ and $t = 2p \cdot \eta$ we deduce that

$$\frac{1}{2} (|p + \eta|^{2s} + |p - \eta|^{2s}) - |p|^{2s} \leq (|p|^2 + |\eta|^2)^s - |p|^{2s}.$$

Next, for $0 < s < 1$ concavity implies that $(a+b)^s \leq a^s + sa^{s-1}b$ for $a, b > 0$, from which we learn that

$$(|p|^2 + |\eta|^2)^s - |p|^{2s} \leq s|p|^{2(s-1)}|\eta|^2.$$

Hence, replacing η with $h\eta$ and using (14) we can estimate

$$R_h(p) \leq s \int |p|^{-2+2s} |h\eta|^2 |\hat{\phi}(\eta)|^2 d\eta = s |p|^{-2+2s} h^2 \int |\nabla\phi|^2 dx \leq Ch^2 |p|^{-2+2s}.$$

Thus the upper bound follows from (21) and (25).

3. ASYMPTOTICS ON THE HALF-SPACE

Our goal in this section is to prove the analogue of Proposition 5 in the case where Ω is the half-space $\mathbb{R}_+^d = \{(x', x_d) : x_d > 0\}$. We define the operator H^+ on $L^2(\mathbb{R}_+^d)$, in the same way as H_Ω , with form domain

$$\mathcal{H}^s(\mathbb{R}_+^d) = \left\{ v \in H^s(\mathbb{R}^d) : v \equiv 0 \text{ on } \mathbb{R}^d \setminus \overline{\mathbb{R}_+^d} \right\}.$$

We shall prove

Proposition 8. *Assume that $\phi \in C_0^1(\mathbb{R}^d)$ is supported in a ball of radius $l > 0$ and assume that (14) is satisfied. Then for $h > 0$ and any $0 < \delta_1 < 1$ and $0 < \delta_2 < \min\{1, 2s\}$ we have*

$$\begin{aligned} & -C_{\delta_1, \delta_2} (l^{d-1-\delta_1} h^{-d+1+\delta_1} + l^{d-1-\delta_2} h^{-d+1+\delta_2}) \\ & \leq \text{Tr}(\phi H^+ \phi)_- - L_{s,d}^{(1)} \int_{\mathbb{R}_+^d} \phi^2(x) dx h^{-d} + L_{s,d}^{(2)} \int_{\mathbb{R}^{d-1}} \phi^2(x', 0) dx' h^{-d+1} \\ & \leq C_{\delta_1} l^{d-1-\delta_1} h^{-d+1+\delta_1}. \end{aligned}$$

This result depends on a more or less explicit diagonalization of the operator H^+ , which is far from obvious. This is accomplished in Subsections 3.1 and 3.2, relying crucially on recent results of Kwaśnicki [Kwa10b] about non-local operators on a half-line. These results are collected and extended to our needs in Section 8.

3.1. The model operator on the half-line. In this subsection we collect some facts about the one-dimensional operator

$$A^+ = \left(-\frac{d^2}{dt^2} + 1 \right)^s$$

in $L^2(\mathbb{R}_+)$ with form domain $\mathcal{H}^s(\mathbb{R}_+)$, and about the corresponding operator A in $L^2(\mathbb{R})$, defined analogously to A^+ , but with form domain $H^s(\mathbb{R})$.

For $\mu > 0$ and $t, u \in \mathbb{R}_+$, let $e^+(t, u, \mu)$ and $a^+(t, u, \mu)$ be the integral kernels of $(A^+ - \mu)_-^0$ and $(A^+ - \mu)_-$, respectively. Similarly, we define $a(t, u, \mu)$ via $(A - \mu)_-$. To simplify notation we abbreviate $a^+(t, \mu) = a^+(t, t, \mu)$. We also note that $a(\mu) = a(t, t, \mu)$ is independent of $t \in \mathbb{R}_+$. The inequality $A^+ \geq 1$ implies that $a^+(t, u, \mu) = e^+(t, u, \mu) = 0$ for $\mu < 1$ and similarly for $a(t, u, \mu)$ and $e(t, u, \mu)$.

The following two results about $e^+(t, \mu)$ and $a^+(t, \mu)$ are rather technical and we defer the proofs to Subsections 8.1 and 8.2. The first one provides a rough a-priori bound on $e^+(t, u, \mu)$.

Lemma 9. *For any $\mu > 0$ and $t, u \in \mathbb{R}_+$ one has $|e^+(t, u, \mu)| \leq C\mu^{1/2s}$.*

The second result in this subsection quantifies that $a^+(t, \mu)$ is close to $a(\mu)$ for large t .

Lemma 10. *For any $0 \leq \gamma < 1$ there is a constant C_γ such that for all $\mu \geq 1$,*

$$\int_0^\infty t^\gamma |a^+(t, \mu) - a(\mu)| dt \leq C_\gamma \mu ((\ln \mu)^2 + 1). \quad (26)$$

In particular, the function

$$K(t) = \frac{1}{(2\pi)^{d-1}} \int_{\mathbb{R}^{d-1}} |\xi'|^{1+2s} (a(|\xi'|^{-2s}) - a^+(t|\xi'|, |\xi'|^{-2s})) d\xi', \quad t > 0, \quad (27)$$

satisfies for every $0 \leq \gamma < 1$

$$\int_0^\infty t^\gamma |K(t)| dt < \infty.$$

With this lemma at hand we can now define the constant $L_{s,d}^{(2)}$ which appears in our main theorem by

$$L_{s,d}^{(2)} = \int_0^\infty K(t) dt. \quad (28)$$

(This integral converges by Lemma 10.) Expression (28) suffices for the proof of our main result. In Section 6, see also (71), we will derive different representation for $L_{s,d}^{(2)}$.

3.2. Reduction from the half-space to the half-line. Our goal in this subsection is to write the spectral projections of the operator H^+ on the half-space in terms of those of the operator A^+ on the half-line. Before turning to spectral projections we treat resolvents.

Lemma 11. *For $x = (x', x_d) \in \mathbb{R}_+^d$, $y = (y', y_d) \in \mathbb{R}_+^d$ and $z \in \mathbb{C} \setminus [0, \infty)$ the resolvent kernels of H^+ and of A^+ are related by*

$$\begin{aligned} & (H^+ - z)^{-1}(x, y) \\ &= \frac{1}{h^d} \int_{\mathbb{R}^{d-1}} |\xi'|^{1-2s} e^{i\xi' \cdot (x' - y')/h} \left(A^+ - \frac{z}{|\xi'|^{2s}} \right)^{-1} \left(\frac{x_d |\xi'|}{h}, \frac{y_d |\xi'|}{h} \right) \frac{d\xi'}{(2\pi)^{d-1}}. \end{aligned}$$

This lemma, together with the representations (see, e.g., [Kat66])

$$(H^+)_-^0 = -\frac{1}{2\pi i} \int_{\Gamma(h^{-2s})} (H^+ - z)^{-1} dz$$

and

$$(H^+)_- = -h^{2s} \int_0^{h^{-2s}} \frac{1}{2\pi i} \int_{\Gamma(\nu)} (H^+ - z)^{-1} dz d\nu,$$

where $\Gamma(\nu) = \{z \in \mathbb{C} : |z| = \nu\}$, implies that

$$(H^+)_-^0(x, y) = \frac{1}{h^d} \int_{\mathbb{R}^{d-1}} |\xi'| e^{i\xi' \cdot (x' - y')/h} e^+ \left(\frac{x_d |\xi'|}{h}, \frac{y_d |\xi'|}{h}, \frac{1}{|\xi'|^{2s}} \right) \frac{d\xi'}{(2\pi)^{d-1}} \quad (29)$$

and

$$(H^+)_-(x, y) = \frac{1}{h^d} \int_{\mathbb{R}^{d-1}} |\xi'|^{1+2s} e^{i\xi' \cdot (x' - y')/h} a^+ \left(\frac{x_d |\xi'|}{h}, \frac{y_d |\xi'|}{h}, \frac{1}{|\xi'|^{2s}} \right) \frac{d\xi'}{(2\pi)^{d-1}}. \quad (30)$$

We now give the

Proof of Lemma 11. By scaling we may assume that $h = 1$. Given $f \in L^2(\mathbb{R}_+^d)$ we want to solve $(-\Delta)^s u = zu + f$. Take $\psi \in C_0^\infty(\mathbb{R}^{d-1})$ and $\phi \in C_0^\infty(0, \infty)$. Then $\psi \otimes \phi$ belongs to the form domain of H^+ and therefore the equation implies that

$$\begin{aligned} \iint (|\xi'|^2 + \xi_d^2)^s \overline{\hat{\psi}(\xi') \hat{\phi}(\xi_d)} \hat{u}(\xi', \xi_d) d\xi' d\xi_d &= z \iint \overline{\hat{\psi}(\xi') \hat{\phi}(\xi_d)} \hat{u}(\xi', \xi_d) d\xi' d\xi_d \\ &+ \iint \overline{\hat{\psi}(\xi') \hat{\phi}(\xi_d)} \hat{f}(\xi', \xi_d) d\xi' d\xi_d. \end{aligned}$$

Since ψ is arbitrary, this means that for a.e. ξ' ,

$$\int (|\xi'|^2 + \xi_d^2)^s \overline{\hat{\phi}(\xi_d)} \hat{u}(\xi', \xi_d) d\xi_d = z \int \overline{\hat{\phi}(\xi_d)} \hat{u}(\xi', \xi_d) d\xi_d + \int \overline{\hat{\phi}(\xi_d)} \hat{f}(\xi', \xi_d) d\xi_d. \quad (31)$$

For fixed ξ' define functions $v_{\xi'}$, $g_{\xi'}$ and $\chi_{\xi'}$ on $(0, \infty)$ by

$$\begin{aligned} v_{\xi'}(t) &:= (2\pi)^{-(d-1)/2} \int_{\mathbb{R}^{d-1}} |\xi'|^{-1} u(x', |\xi'|^{-1}t) e^{-i\xi' \cdot x'} dx', \\ g_{\xi'}(t) &:= (2\pi)^{-(d-1)/2} \int_{\mathbb{R}^{d-1}} |\xi'|^{-1-2s} f(x', |\xi'|^{-1}t) e^{-i\xi' \cdot x'} dx', \\ \chi_{\xi'}(t) &:= |\xi'|^{-1} \phi(|\xi'|^{-1}t). \end{aligned}$$

We note that the one-dimensional Fourier transforms of these functions are given by

$$\begin{aligned} \widehat{v_{\xi'}}(k) &= \hat{u}(\xi', |\xi'|k), \\ \widehat{g_{\xi'}}(k) &= |\xi'|^{-2s} \hat{f}(\xi', |\xi'|k), \\ \widehat{\chi_{\xi'}}(k) &= \hat{\phi}(|\xi'|k). \end{aligned}$$

Hence we can rewrite (31) as

$$\int (1 + k^2)^s \overline{\widehat{\chi_{\xi'}}(k)} \widehat{v_{\xi'}}(k) dk = |\xi'|^{-2s} z \int \overline{\widehat{\chi_{\xi'}}(k)} \widehat{v_{\xi'}}(k) dk + \int \overline{\widehat{\chi_{\xi'}}(k)} \widehat{g_{\xi'}}(k) dk.$$

Note that both $v_{\xi'}$ and $\chi_{\xi'}$ belong to the form domain of A^+ and that the set of all functions $\chi_{\xi'}$ obtained in this way is dense in the form sense (by the definition of A^+). Therefore the equation can be written as

$$A^+ v_{\xi'} = |\xi'|^{-2s} z v_{\xi'} + g_{\xi'}.$$

We abbreviate $r_z^+(t, w) = (A^+ - z)^{-1}(t, w)$ and conclude that

$$v_{\xi'}(t) = \int_0^\infty r_{z|\xi'|^{-2s}}^+(t, w) g_{\xi'}(w) dw.$$

Recalling the definitions of $v_{\xi'}$ and $g_{\xi'}$ this reads

$$\begin{aligned} & \int_{\mathbb{R}^{d-1}} |\xi'|^{-1} u(x', |\xi'|^{-1} t) e^{-i\xi' \cdot x'} dx' \\ &= \int_0^\infty \int_{\mathbb{R}^{d-1}} r_{z|\xi'|^{-2s}}^+(t, w) |\xi'|^{-1-2s} f(x', |\xi'|^{-1} t) e^{-i\xi' \cdot x'} dw dx'. \end{aligned}$$

Multiplying by $|\xi'|$, setting $t = |\xi'| x_d$ and inverting the Fourier transform, we obtain

$$\begin{aligned} u(x) &= (2\pi)^{-d+1} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}^{d-1}} \int_0^\infty r_{z|\xi'|^{-2s}}^+(|\xi'| x_d, w) |\xi'|^{-2s} f(y', |\xi'|^{-1} w) e^{i\xi' \cdot (x' - y')} dw d\xi' dy' \\ &= (2\pi)^{-d+1} \int_{\mathbb{R}^d} \int_{\mathbb{R}^{d-1}} r_{z|\xi'|^{-2s}}^+(|\xi'| x_d, |\xi'| y_d) |\xi'|^{1-2s} f(y) e^{i\xi' \cdot (x' - y')} d\xi' dy. \end{aligned}$$

This proves the lemma. \square

3.3. Proof of Proposition 8. Our next step is to state upper and lower bounds on $\text{Tr}(\phi H^+ \phi)_-$ in terms of the one-dimensional model operators A and A^+ , in particular, in terms of the function $K(t)$ given in (27). As explained below, the main result of this section, Proposition 8, will be a direct consequence of the following estimates.

Proposition 12. *Assume that $\phi \in C_0^1(\mathbb{R}^d)$ is supported in a ball of radius $l = 1$ and assume that (14) is satisfied with $l = 1$. Then for any $0 < \delta_2 < \min\{1, 2s\}$ there is a constant C_{δ_2} such that for all $h > 0$ we have*

$$\text{Tr}(\phi H^+ \phi)_- \leq L_{s,d}^{(1)} \int_{\mathbb{R}_+^d} \phi^2(x) dx h^{-d} - \int_{\mathbb{R}_+^d} \phi^2(x) \frac{1}{h} K\left(\frac{x_d}{h}\right) dx h^{-d+1}, \quad (32)$$

$$\text{Tr}(\phi H^+ \phi)_- \geq L_{s,d}^{(1)} \int_{\mathbb{R}_+^d} \phi^2(x) dx h^{-d} - \int_{\mathbb{R}_+^d} \phi^2(x) \frac{1}{h} K\left(\frac{x_d}{h}\right) dx h^{-d+1} - C_{\delta_2} h^{-d+1+\delta_2}. \quad (33)$$

Assuming Proposition 12, we now give the short

Proof of Proposition 8. To prove the proposition we may rescale ϕ and hence assume $l = 1$. Proposition 8 is then an immediate consequence of Proposition 12 provided we can show that for any $0 < \delta_1 < 1$ there is a C_{δ_1} such that for all $h > 0$

$$\left| \int_{\mathbb{R}^{d-1}} \phi^2(x) \frac{1}{h} K\left(\frac{x_d}{h}\right) dx - L_{s,d}^{(2)} \int_{\mathbb{R}^{d-1}} \phi^2(x', 0) dx' \right| \leq C_{\delta_1} h^{\delta_1}. \quad (34)$$

In order to obtain the latter bound, we substitute $x_d = th$ and write, recalling (28),

$$\begin{aligned} & \int_{\mathbb{R}^{d-1}} \phi^2(x) \frac{1}{h} K\left(\frac{x_d}{h}\right) dx - L_{s,d}^{(2)} \int_{\mathbb{R}^{d-1}} \phi^2(x', 0) dx' \\ &= \int_0^\infty K(t) \int_{\mathbb{R}^{d-1}} \int_0^{th} \partial_\tau \phi^2(x', \tau) d\tau dx' dt. \end{aligned}$$

By Hölder's inequality we can further estimate

$$\begin{aligned} \left| \int_{\mathbb{R}^{d-1}} \int_0^{th} \partial_\tau \phi^2(x', \tau) d\tau dx' \right| &\leq \left(\int_0^{th} d\tau \right)^{\delta_1} \left(\int_0^\infty \left| \int_{\mathbb{R}^{d-1}} \partial_\tau \phi^2(x', \tau) dx' \right|^{(1-\delta_1)^{-1}} d\tau \right)^{1-\delta_1} \\ &\leq Ct^{\delta_1} h^{\delta_1}. \end{aligned}$$

Since $\int_0^\infty t^{\delta_1} |K(t)| dt < \infty$ by Lemma 10, we obtain inequality (34). \square

In the following two subsections we shall prove the lower and the upper bound in Proposition 12, respectively.

3.4. Lower bound. To prove (32) we use that

$$-\mathrm{Tr}(\phi H^+ \phi)_- \geq -\mathrm{Tr}(\phi(H^+)_- \phi).$$

The lower bound follows from this by integrating the identity

$$(H^+)_-(x, x) = h^{-d} L_{s,d}^{(1)} - h^{-d} K\left(\frac{x_d}{h}\right), \quad (35)$$

against ϕ^2 . Equation (35) is a consequence of (30). Indeed, by the same argument as in Subsection 3.2 we learn that

$$(H_0)_-(x, x) = \frac{1}{(2\pi)^{d-1}} \frac{1}{h^d} \int_{\mathbb{R}^{d-1}} |\xi'|^{1+2s} a(|\xi'|^{-2s}) d\xi'.$$

On the other hand, by direct diagonalization as in Subsection 2.1 we find that

$$(H_0)_-(x, x) = h^{-d} L_{s,d}^{(1)}.$$

Comparing these two identities with (30) we arrive at (35), thus establishing (32).

3.5. Upper bound. To prove (33) we set $\gamma = (H^+)_-$. Its integral kernel is given by (29) in terms of the kernel $e^+(\cdot, \cdot, \mu)$ of $(A^+ - \mu)_-$. By the variational principle it follows that

$$\begin{aligned} -\mathrm{Tr}(\phi H^+ \phi)_- &\leq \mathrm{Tr}(\phi \gamma \phi H^+) \\ &= \frac{1}{h^{2d}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}^d} |\xi'| e^{i\xi' \cdot (x' - y')/h} e^+(x_d |\xi'| h^{-1}, y_d |\xi'| h^{-1}, |\xi'|^{-2s}) \\ &\quad \times (|p|^{2s} - 1) e^{ip \cdot (y-x)/h} \phi(x) \phi(y) \frac{dp d\xi' dx dy}{(2\pi)^{2d-1}}. \end{aligned} \quad (36)$$

We insert the identity

$$\phi(x) \phi(y) = \frac{1}{2} (\phi^2(x) + \phi^2(y) - |\phi(x) - \phi(y)|^2),$$

use the symmetry in x and y and substitute $q = p_d/|p'|$ to obtain

$$-\mathrm{Tr}(\phi H^+ \phi) \leq I_h[\phi] - R_h[\phi]$$

with the main term

$$I_h[\phi] = \frac{1}{h^{2d}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} |\xi'| e^{i(\xi' - p') \cdot (x' - y')/h} e^+ (x_d |\xi'| h^{-1}, y_d |\xi'| h^{-1}, |\xi'|^{-2s}) \\ \times e^{i(y_d - x_d)|p'|q/h} ((q^2 + 1)^s - |p'|^{-2s}) |p'|^{1+2s} \phi^2(x) \frac{dq dp' d\xi' dx dy}{(2\pi)^{2d-1}}$$

and the remainder

$$R_h[\phi] = \frac{1}{h^{2d}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}^d} |\xi'| e^{i\xi' \cdot (x' - y')/h} e^+ (x_d |\xi'| h^{-1}, y_d |\xi'| h^{-1}, |\xi'|^{-2s}) \\ \times |p|^{2s} e^{ip \cdot (y - x)/h} |\phi(x) - \phi(y)|^2 \frac{dp d\xi' dx dy}{2(2\pi)^{2d-1}}.$$

Since $\phi \in C_0^1(\mathbb{R}^d)$ we can perform the y' -integration in $I_h[\phi]$. We use the fact that

$$\int_{\mathbb{R}} \int_0^\infty e^+ (x_d, y_d, \mu) ((q^2 + 1)^s - \mu) e^{i(y_d - z_d)q} dy_d dq = -a^+ (x_d, z_d, \mu)$$

and obtain

$$I_h[\phi] = \frac{1}{h^{d+1}} \int_{\mathbb{R}_+^d} \int_0^\infty \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} |\xi'|^{2s+2} e^+ (x_d |\xi'| h^{-1}, y_d |\xi'| h^{-1}, |\xi'|^{-2s}) \\ \times ((q^2 + 1)^s - |\xi'|^{-2s}) e^{i(y_d - x_d)|\xi'|q/h} \phi^2(x) \frac{dq d\xi' dy_d dx}{(2\pi)^d} \\ = -\frac{1}{h^d} \int_{\mathbb{R}_+^d} \phi^2(x) \int_{\mathbb{R}^{d-1}} |\xi'|^{2s+1} a^+ (x_d |\xi'| h^{-1}, |\xi'|^{-2s}) \frac{d\xi' dx}{(2\pi)^{d-1}}.$$

Using again (35) we find that

$$I_h[\phi] = -L_{s,d}^{(1)} \int_{\mathbb{R}_+^d} \phi^2(x) dx h^{-d} + \int_{\mathbb{R}_+^d} \phi^2(x) K\left(\frac{x_d}{h}\right) dx h^{-d}. \quad (37)$$

It remains to study $R_h[\phi]$. We claim that for any $\frac{1}{2} - s < \sigma < \min\{\frac{1}{2}, 1 - s\}$ there is a C_σ such that

$$|R_h[\phi]| \leq C_\sigma h^{-d+2s+2\sigma} \quad (38)$$

for all $h > 0$. This, together with (37) will complete the proof of (33).

In order to show (38) we perform the p integration and find that

$$R_h[\phi] = -\frac{C}{h^{d-2s}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}^{d-1}} |\xi'| e^{i\xi' \cdot (x' - y')/h} e^+ \left(\frac{x_d |\xi'|}{h}, \frac{y_d |\xi'|}{h}, \frac{1}{|\xi'|^{2s}} \right) \\ \times \frac{|\phi(x) - \phi(y)|^2}{|x - y|^{d+2s}} d\xi' dx dy.$$

We insert

$$e^{i\xi' \cdot (x' - y')/h} = \frac{h^{2\sigma}}{|\xi'|^{2\sigma}} (-\Delta_{x'})^\sigma e^{i\xi' \cdot (x' - y')/h}$$

and integrate by parts to get

$$R_h[\phi] = -\frac{C}{h^{d-2s-2\sigma}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}^{d-1}} |\xi'|^{1-2\sigma} e^{i\xi' \cdot (x'-y')/h} e^+ \left(\frac{x_d |\xi'|}{h}, \frac{y_d |\xi'|}{h}, \frac{1}{|\xi'|^{2s}} \right) d\xi' \\ \times (-\Delta_{x'})^\sigma \frac{|\phi(x) - \phi(y)|^2}{|x-y|^{d+2s}} dx dy.$$

By Lemma 9 and the fact that $e^+(t, u, \mu) = 0$ for $\mu \leq 1$ we arrive at

$$|R_h[\phi]| \leq \frac{C}{h^{d-2s-2\sigma}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \int_{\{\xi' \in \mathbb{R}^{d-1}; |\xi'| < 1\}} |\xi'|^{-2\sigma} d\xi' \left| (-\Delta_{x'})^\sigma \frac{|\phi(x) - \phi(y)|^2}{|x-y|^{d+2s}} \right| dx dy \\ \leq \frac{C}{h^{d-2s-2\sigma}} \int_{\mathbb{R}_+^d} \int_{\mathbb{R}_+^d} \left| (-\Delta_{x'})^\sigma \frac{|\phi(x) - \phi(y)|^2}{|x-y|^{d+2s}} \right| dx dy.$$

According to Lemma 24 this implies (38) and hence completes the proof of (33).

4. LOCAL ASYMPTOTICS NEAR THE BOUNDARY

In this section we prove Proposition 5. After having analyzed the half-space case in the previous section, we now show how the case of a general domain follows. We shall transform the operator H_Ω locally to an operator given on the half-space $\mathbb{R}_+^d = \{(y', y_d) \in \mathbb{R}^{d-1} \times \mathbb{R} : y_d > 0\}$ and we shall quantify the error made by this straightening of the boundary.

Under the conditions of Proposition 5, let B denote the open ball of radius $l > 0$, containing the support of ϕ . For $x_0 \in B \cap \partial\Omega$ let ν_{x_0} be the inner normal unit vector at x_0 . We choose a Cartesian coordinate system such that $x_0 = 0$ and $\nu_{x_0} = (0, \dots, 0, 1)$, and we write $x = (x', x_d) \in \mathbb{R}^{d-1} \times \mathbb{R}$ for $x \in \mathbb{R}^d$.

For sufficiently small $l > 0$ one can introduce new local coordinates near the boundary. Let D denote the projection of B on the hyperplane given by $x_d = 0$. Since the boundary of Ω is compact and $C^{1,\alpha}$ there is a constant $c > 0$ such that for $0 < l \leq c$ we can find a real function $f \in C^{1,\alpha}$ given on D , satisfying

$$\partial\Omega \cap B = \{(x', x_d) : x' \in D, x_d = f(x')\} \cap B.$$

The choice of coordinates implies $f(0) = 0$ and $\nabla f(0) = 0$. Hence, we can estimate

$$\sup_{x' \in D} |\nabla f(x')| = \sup_{x' \in D} |\nabla f(x') - \nabla f(0)| \leq C_f |x'|^\alpha \leq C_f l^\alpha.$$

Since the boundary of Ω is compact we can choose a constant $C > 0$, depending only on Ω , in particular independent of f , such that the bound

$$\sup_{x' \in D} |\nabla f(x')| \leq Cl^\alpha \tag{39}$$

holds.

We introduce new local coordinates via the diffeomorphism $\varphi : D \times \mathbb{R} \rightarrow \mathbb{R}^d$, given by

$$y_j = \varphi_j(x) = x_j \quad \text{for } j = 1, \dots, d-1$$

and

$$y_d = \varphi_d(x) = x_d - f(x').$$

Note that the determinant of the Jacobian matrix of φ equals 1 and that the inverse of φ is given on $\text{ran } \varphi = D \times \mathbb{R}$. In particular, we get

$$\varphi(\partial\Omega \cap B) \subset \partial\mathbb{R}_+^d = \{y \in \mathbb{R}^d : y_d = 0\}.$$

Fix $v \in \mathcal{H}^s(\Omega)$ with support in \overline{B} . For $y \in \text{ran } \varphi$ put $\tilde{v}(y) = v \circ \varphi^{-1}(y)$ and extend \tilde{v} by zero to \mathbb{R}^d .

Lemma 13. *The function \tilde{v} belongs to $\mathcal{H}^s(\mathbb{R}_+^d)$ and for $0 < l \leq c$ we have*

$$\left| (\tilde{v}, (-\Delta)_{\mathbb{R}_+^d}^s \tilde{v}) - (v, (-\Delta)_{\Omega}^s v) \right| \leq C l^\alpha \min \left\{ (\tilde{v}, (-\Delta)_{\mathbb{R}_+^d}^s \tilde{v}), (v, (-\Delta)_{\Omega}^s v) \right\}.$$

Proof. By definition, \tilde{v} belongs to $H^s(\mathbb{R}^d)$ and for $y \in \mathbb{R}^d \setminus \overline{\mathbb{R}_+^d}$ we find $x_d = y_d + f(y') < f(x')$, thus $\tilde{v}(y) = v(x) = 0$. Therefore \tilde{v} belongs to $\mathcal{H}^s(\mathbb{R}_+^d)$.

Using the new local coordinates we get

$$(v, (-\Delta)_{\Omega}^s v) = C_{s,d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|v(x) - v(w)|^2}{|x - w|^{d+2s}} dx dw = C_{s,d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|\tilde{v}(y) - \tilde{v}(z)|^2}{|x - w|^{d+2s}} dy dz, \quad (40)$$

where $y = \varphi(x)$ and $z = \varphi(w)$, thus $x = (y', y_d + f(y'))$ and $w = (z', z_d + f(z'))$. Let us write

$$\begin{aligned} & \left| \frac{1}{|y - z|^{d+2s}} - \frac{1}{|x - w|^{d+2s}} \right| \\ &= \frac{1}{|y - z|^{d+2s}} \left| 1 - \frac{|y - z|^{d+2s}}{[|y' - z'|^2 + (y_d + f(y') - z_d - f(z'))^2]^{d/2+s}} \right|. \end{aligned}$$

After multiplying out, the last fraction equals

$$\left(1 + \frac{(f(y') - f(z'))^2 + 2(y_d - z_d)(f(y') - f(z'))}{|y - z|^2} \right)^{-(d/2+s)}$$

and we can employ (39) to estimate

$$\begin{aligned} & \left| \frac{(f(y') - f(z'))^2 + 2(y_d - z_d)(f(y') - f(z'))}{|y - z|^2} \right| \\ & \leq \sup |\nabla f|^2 \frac{|y' - z'|^2}{|y - z|^2} + 2 \sup |\nabla f| \frac{|y' - z'| |y_d - z_d|}{|y - z|^2} \leq C l^\alpha. \end{aligned}$$

Choosing l small enough we can assume $C l^\alpha < 1/2$. Then, combining the foregoing relations, we find

$$\left| \frac{1}{|x - w|^{d+2s}} - \frac{1}{|y - z|^{d+2s}} \right| \leq C \frac{l^\alpha}{|y - z|^{d+2s}}. \quad (41)$$

From (40) and (41) we conclude

$$\begin{aligned} & \left| (\tilde{v}, (-\Delta)_{\mathbb{R}_+^d}^s \tilde{v}) - (v, (-\Delta)_\Omega^s v) \right| \\ & \leq C_{s,d} \iint |\tilde{v}(y) - \tilde{v}(z)|^2 \left| \frac{1}{|y-z|^{d+2s}} - \frac{1}{|x-w|^{d+2s}} \right| dy dz \\ & \leq C l^\alpha (\tilde{v}, (-\Delta)_{\mathbb{R}_+^d}^s \tilde{v}). \end{aligned}$$

This proves the first claim of the Lemma. The second claim follows by interchanging the roles of $(-\Delta)_{\mathbb{R}_+^d}^s$ and $(-\Delta)_\Omega^s$. \square

On the range of φ we define $\tilde{\phi}_u = \phi_u \circ \varphi^{-1}$ and extend it by zero to \mathbb{R}^d such that $\tilde{\phi}_u \in C_0^1(\mathbb{R}^d)$ and $\|\nabla \tilde{\phi}_u\|_\infty \leq C l^{-1}$ hold. Using Lemma 13 we show the following relations.

Lemma 14. *For $0 < l \leq c$ and any $h > 0$ the estimate*

$$\left| \text{Tr}(\phi H_\Omega \phi)_- - \text{Tr}(\tilde{\phi} H^+ \tilde{\phi})_- \right| \leq C l^{d+\alpha} h^{-d} \quad (42)$$

holds. Moreover, we have

$$\int_\Omega \phi^2(x) dx = \int_{\mathbb{R}_+^d} \tilde{\phi}^2(y) dy \quad (43)$$

and

$$0 \leq \int_{\partial\Omega} \phi^2(x) d\sigma(x) - \int_{\mathbb{R}^{d-1}} \tilde{\phi}^2(y', 0) dy' \leq C l^{d-1+2\alpha}. \quad (44)$$

Proof. The definition of $\tilde{\phi}$ and the fact that the Jacobian of ϕ equals 1 immediately gives (43). Using (39) we estimate

$$\int_{\partial\Omega} \phi^2(x) d\sigma(x) = \int_{\mathbb{R}^{d-1}} \tilde{\phi}^2(y', 0) \sqrt{1 + |\nabla f|^2} dy' \leq \int_{\mathbb{R}^{d-1}} \tilde{\phi}^2(y', 0) dy' + C l^{d-1+2\alpha}.$$

from which (44) follows.

To prove (42) we refer to the variational principle once more and note that

$$-\text{Tr}(\phi H_\Omega \phi)_- = \inf_{0 \leq \gamma \leq 1} \text{Tr}(\phi \gamma \phi H_\Omega),$$

where we can assume that infimum is taken over trial density matrices γ supported in $\overline{B} \times \overline{B}$. Fix such a γ . For y and z from $D \times \mathbb{R}$ set

$$\tilde{\gamma}(y, z) = \gamma(\varphi^{-1}(y), \varphi^{-1}(z)),$$

so that $0 \leq \tilde{\gamma} \leq 1$ and the range of $\tilde{\gamma}$ belongs to the form domain of $\tilde{\phi} H^+ \tilde{\phi}$. According to Lemma 13 it follows that

$$\begin{aligned} \text{Tr}(\phi \gamma \phi H_\Omega) & \geq \text{Tr} \left(\tilde{\phi} \tilde{\gamma} \tilde{\phi} \left(h^{2s} (1 - C l^\alpha) (-\Delta)_{\mathbb{R}_+^d}^s - 1 \right) \right) \\ & \geq -\text{Tr} \left(\tilde{\phi} \left((1 - C l^\alpha) h^{2s} (-\Delta)_{\mathbb{R}_+^d}^s - 1 \right) \tilde{\phi} \right)_- \end{aligned}$$

and consequently

$$\mathrm{Tr}(\phi H_\Omega \phi)_- \leq \mathrm{Tr} \left(\tilde{\phi} \left((1 - Cl^\alpha) h^{2s} (-\Delta)_{\mathbb{R}_+^d}^s - 1 \right) \tilde{\phi} \right)_-.$$

Set $\varepsilon = 2Cl^\alpha$ and assume l to be sufficiently small, so that $0 < \varepsilon \leq 1/2$. Then

$$\begin{aligned} \mathrm{Tr}(\phi H_\Omega \phi)_- &\leq \mathrm{Tr} \left(\tilde{\phi} \left((1 - Cl^\alpha) h^{2s} (-\Delta)_{\mathbb{R}_+^d}^s - 1 \right) \tilde{\phi} \right)_- \\ &\leq \mathrm{Tr} \left(\tilde{\phi} \left((-h^2 \Delta)_{\mathbb{R}_+^d}^s - 1 \right) \tilde{\phi} \right)_- + \mathrm{Tr} \left(\tilde{\phi} \left((\varepsilon - Cl^\alpha) h^{2s} (-\Delta)_{\mathbb{R}_+^d}^s - \varepsilon \right) \tilde{\phi} \right)_- \\ &\leq \mathrm{Tr}(\tilde{\phi} H^+ \tilde{\phi})_- + \varepsilon \mathrm{Tr} \left(\tilde{\phi} \left((h^{2s}/2) (-\Delta)_{\mathbb{R}_+^d}^s - 1 \right) \tilde{\phi} \right)_-. \end{aligned}$$

Using Lemma 6 we estimate $\mathrm{Tr}(\tilde{\phi}((h^{2s}/2)(-\Delta)_{\mathbb{R}_+^d}^s - 1)\tilde{\phi})_- \leq Cl^d h^{-d}$ and it follows that

$$\mathrm{Tr}(\phi H_\Omega \phi)_- \leq \mathrm{Tr}(\tilde{\phi} H^+ \tilde{\phi})_- + Cl^{d+\alpha} h^{-d}.$$

Finally, by interchanging the roles of H_Ω and H^+ , we get an analogous lower bound and the proof of the Lemma is complete. \square

We conclude this section by giving the short

Proof of Proposition 5. It suffices to combine Lemma 14 and Proposition 8. \square

5. LOCALIZATION

In this section we construct the family of localization functions $(\phi_u)_{u \in \mathbb{R}^d}$ and prove Proposition 3. Fix a real-valued function $\phi \in C_0^\infty(\mathbb{R}^d)$ with support in the ball $\{x \in \mathbb{R}^d : |x| < 1\}$ that satisfies $\|\phi\|_2 = 1$. We recall the definition of the local length scale $l(u)$ from (11). For $u, x \in \mathbb{R}^d$ let $J(x, u)$ be the Jacobian of the map $u \mapsto (x - u)/l(u)$. We define

$$\phi_u(x) = \phi \left(\frac{x - u}{l(u)} \right) \sqrt{J(x, u)} l(u)^{d/2},$$

such that ϕ_u is supported in the ball $B_u = \{x \in \mathbb{R}^d : |x - u| < l(u)\}$.

By definition, the function $l(u)$ is smooth and satisfies $0 < l(u) \leq 1/2$ and $\|\nabla l\|_\infty \leq 1/2$. Therefore, according to [SS03], the functions ϕ_u satisfy (12) and (13) for all $u \in \mathbb{R}^d$.

To prove the lower bound in Proposition 3 we follow some ideas from [LY88]. In particular, we need the following auxiliary results; the first one gives an IMS-type localization formula for the fractional Laplacian.

Lemma 15. *For the family of functions $(\phi_u)_{u \in \mathbb{R}^d}$ introduced above and for all $f \in \mathcal{H}^s(\Omega)$ the identity*

$$(f, (-\Delta)^s f) = \int_{\Omega^*} (\phi_u f, (-\Delta)^s \phi_u f) l(u)^{-d} du - (f, Lf)$$

holds with $\Omega^* = \{u \in \mathbb{R}^d : \mathrm{supp} \phi_u \cap \Omega \neq \emptyset\}$. The operator L is of the form

$$L = \int_{\Omega^*} L_{\phi_u} l(u)^{-d} du, \quad (45)$$

where L_{ϕ_u} is a bounded operator with integral kernel

$$L_{\phi_u}(x, y) = C_{s,d} \frac{|\phi_u(x) - \phi_u(y)|^2}{|x - y|^{d+2s}} \chi_\Omega(x) \chi_\Omega(y).$$

Here χ_Ω denotes the characteristic function of Ω .

Lemma 15 implies that for any operator γ with range in $\mathcal{H}^s(\Omega)$

$$\mathrm{Tr} \gamma(-\Delta)^s = \int_{\mathbb{R}^d} \mathrm{Tr} (\gamma \phi_u(-\Delta)^s \phi_u) l(u)^{-d} du - \mathrm{Tr} \gamma L. \quad (46)$$

The next result allows to estimate the localization error $\mathrm{Tr} \gamma L$.

Lemma 16. *For $u \in \mathbb{R}^d$ and $0 < \delta \leq 1/2$ we have*

$$\mathrm{Tr} \gamma L_{\phi_u} \leq \mathrm{Tr} \gamma (C \delta^{2-2s} l(u)^{-2s} \chi_\delta \chi_\Omega) + C \|\gamma\| l(u)^{-2s} \delta^{-d+2-2s} r(\delta)$$

with

$$r(\delta) = \begin{cases} 1 & \text{if } 1 - d/4 < s < 1 \\ |\ln \delta| & \text{if } 0 < s = 1 - d/4 \\ \delta^{d+4s-4} & \text{if } 0 < s < 1 - d/4 \end{cases}.$$

where χ_δ denotes the characteristic function of $\{x \in \mathbb{R}^d : |x - u| < l(u)(1 + \delta)\}$.

Proof. By translation and scaling we can assume that $u = 0$ and $l(u) = 1$, and hence $\phi_u = \phi$. (This rescaling changes Ω , but the bound we are going to prove is independent of the domain and therefore not affected by this dilation.) We set

$$L_\phi^1(x, y) = \begin{cases} L_\phi(x, y) \chi_\delta(x) \chi_\delta(y) & \text{if } |x - y| < \delta \\ 0 & \text{if } |x - y| \geq \delta \end{cases},$$

$L_\phi^0(x, y) = L_\phi(x, y) - L_\phi^1(x, y)$ and $\theta(x) = \int L_\phi^1(x, y) dy$. By a simple adaption of the arguments of [LY88, Thm. 10] we find that for any $\epsilon > 0$

$$\mathrm{Tr} \gamma L_\phi \leq \mathrm{Tr} \gamma (\theta + \epsilon \chi_0) + \frac{\|\gamma\|}{2\epsilon} \mathrm{Tr} (L_\phi^0)^2. \quad (47)$$

It remains to bound θ and $\mathrm{Tr}(L_\phi^0)^2$.

We begin by estimating θ . By definition, for $|x| \geq 1 + \delta$ we have $L_\phi^1(x, y) = 0$ and hence $\theta(x) = 0$, and for $|x| < 1 + \delta$ we get

$$\theta(x) = C_{s,d} \int_{\substack{|x-y| < \delta \\ |y| < 1+\delta}} \frac{(\phi(x) - \phi(y))^2}{|x - y|^{d+2s}} dy \leq C \|\nabla \phi\|_\infty^2 \int_{|x-y| < \delta} \frac{1}{|x - y|^{d+2s-2}} dy.$$

Thus, for all $x \in \mathbb{R}^d$

$$\theta(x) \leq C \delta^{2-2s} \chi_\delta(x). \quad (48)$$

Finally, we estimate $\mathrm{Tr}(L_\phi^0)^2$. The symmetry of $L_\phi^0(x, y)$ implies

$$\mathrm{Tr} (L_\phi^0)^2 \leq C \iint_A \left(\frac{(\phi(x) - \phi(y))^2}{|x - y|^{d+2s}} \right)^2 dx dy$$

where A denotes the set $\{(x, y) \in \mathbb{R}^d \times \mathbb{R}^d : |x| < \min(|y|, 1), |x - y| \geq \delta\}$. Set $A_1 = \{(x, y) \in A : |y| \geq 2\}$ and $A_2 = \{(x, y) \in A : |y| < 2\}$. Then

$$\begin{aligned} \text{Tr} (L_\phi^0)^2 &\leq C \iint_{A_1} \left(\frac{\phi^4(x)}{|x - y|^{2d+4s}} \right) dx dy + C \|\nabla \phi\|_\infty^4 \iint_{A_2} \frac{1}{|x - y|^{2d+4s-4}} dx dy \\ &\leq C r(\delta). \end{aligned} \quad (49)$$

Choosing $\varepsilon = \delta^{2-2s}$ and combining (47) with (48) and (49) yields the claimed result. \square

Proof of Proposition 3. We apply Lemma 16 with a parameter $0 < \delta_u \leq 1/2$ to be specified later. For ease of notation we write χ_u instead of χ_{δ_u} . Identities (45) and (46) and the estimate from Lemma 16 imply

$$\begin{aligned} \text{Tr} \gamma(-\Delta)^s &\geq \int_{\Omega^*} \text{Tr} \gamma \left(\phi_u(-\Delta)^s \phi_u - C \delta_u^{2-2s} l(u)^{-2s} \chi_u(x) \chi_\Omega(x) \right) l(u)^{-d} du \\ &\quad - C \|\gamma\| \int_{\Omega^*} \delta_u^{-d+2-2s} r(\delta_u) l(u)^{-d-2s} du. \end{aligned} \quad (50)$$

If the supports of χ_u and $\phi_{u'}$ overlap, we have $|u - u'| \leq (3/2)l(u) + l(u')$. It follows that $l(u') - l(u) \leq \|\nabla l\|_\infty ((3/2)l(u) + l(u'))$. Since $\|\nabla l\|_\infty \leq 1/2$ we find $l(u') \leq Cl(u)$ and $l(u)^{-1} \leq Cl(u')^{-1}$. Similarly, we get $l(u) \leq Cl(u')$. We assume now that δ_u satisfies

$$\delta_u \leq C \delta_{u'} \quad \text{if } |u - u'| \leq (3/2)(l(u) + l(u')). \quad (51)$$

Using these locally uniform bounds on $l(u)/l(u')$ and $\delta_u/\delta_{u'}$, together with (13), we can deduce the pointwise bound for all $x \in \mathbb{R}^d$

$$\begin{aligned} &\int_{\Omega^*} \delta_u^{2-2s} l(u)^{-2s} \chi_u(x) \chi_\Omega(x) \frac{du}{l(u)^d} \\ &= \int_{\Omega^*} \delta_u^{2-2s} l(u)^{-2s} \chi_u(x) \chi_\Omega(x) \left(\int \phi_{u'}^2(x) \frac{du'}{l(u')^d} \right) \frac{du}{l(u)^d} \\ &\leq C \int_{\Omega^*} \phi_{u'}(x) \delta_{u'}^{2-2s} l(u')^{-2s} \phi_{u'}(x) \frac{du'}{l(u')^d}. \end{aligned}$$

Rewriting the last integral with u as integration variable, in view of (50), we find

$$\begin{aligned} \text{Tr} \gamma(-\Delta)^s &\geq \int_{\Omega^*} \text{Tr} \gamma \left(\phi_u \left((-\Delta)^s - \frac{C \delta_u^{2-2s}}{l(u)^{2s}} \right) \phi_u \right) \frac{du}{l(u)^d} \\ &\quad - C \|\gamma\| \int_{\Omega^*} \delta_u^{-d+2-2s} r(\delta_u) \frac{du}{l(u)^{d+2s}}. \end{aligned}$$

By the variational principle it follows that

$$\begin{aligned} \text{Tr}(H_\Omega)_- &= - \inf_{0 \leq \gamma \leq 1} \text{Tr} \gamma \left((-h^2 \Delta)^s - 1 \right) \\ &\leq \int_{\Omega^*} \text{Tr} \left(\phi_u \left((-h^2 \Delta)^s - 1 - Ch^{2s} \delta_u^{2-2s} l(u)^{-2s} \right) \phi_u \right)_- \frac{du}{l(u)^d} \\ &\quad + Ch^{2s} \int_{\Omega^*} \delta_u^{-d+2-2s} r(\delta_u) \frac{du}{l(u)^{d+2s}}. \end{aligned} \quad (52)$$

To bound the first term, we use Lemma 6. For any $u \in \mathbb{R}^d$, let ρ_u be another parameter satisfying $0 < \rho_u \leq 1/2$ and estimate

$$\begin{aligned} & \text{Tr} \left(\phi_u \left((-h^2 \Delta)^s - 1 - Ch^{2s} \delta_u^{2-2s} l(u)^{-2s} \right) \phi_u \right)_- \\ & \leq \text{Tr} (\phi_u H_\Omega \phi_u)_- + C \text{Tr} \left(\phi_u \left(\rho_u h^{2s} (-\Delta)^s - \rho_u - h^{2s} \delta_u^{2-2s} l(u)^{-2s} \right) \phi_u \right)_- \\ & \leq \text{Tr} (\phi_u H_\Omega \phi_u)_- + Cl(u)^d (\rho_u h^{2s})^{-d/(2s)} (\rho_u + h^{2s} \delta_u^{2-2s} l(u)^{-2s})^{1+d/(2s)}. \end{aligned}$$

We pick $\rho_u = h^{2s} \delta_u^{2-2s} l(u)^{-2s}$. By (17) and our assumption that $\delta_u \leq 1/2$, we see that $\rho_u \leq (h/l_0)^{2s} 2^{6s-2}$. We assume now that $h \leq C^{-1} l_0$ (with a possibly large constant C) in order to guarantee that $\rho_u \leq 1/2$. With this choice we find

$$\text{Tr} \left(\phi_u \left((-h^2 \Delta)^s - 1 - \frac{Ch^{2s} \delta_u^{2-2s}}{l(u)^{2s}} \right) \phi_u \right)_- \leq \text{Tr} (\phi_u H_\Omega \phi_u)_- + C \frac{\delta_u^{2-2s} l(u)^{d-2s}}{h^{d-2s}}. \quad (53)$$

Combining (52) and (53) we obtain

$$\text{Tr}(H_\Omega)_- \leq \int_{\Omega^*} \text{Tr} (\phi_u H_\Omega \phi_u)_- \frac{du}{l(u)^d} + C \int_{\Omega^*} \left(\frac{\delta_u^{2-2s}}{h^{d-2s} l(u)^{2s}} + \frac{h^{2s} \delta_u^{-d+2-2s} r(\delta_u)}{l(u)^{d+2s}} \right) du. \quad (54)$$

At this point we choose δ_u in order to minimize the second integrand, which we shall denote by I_u . We pick

$$\delta_u = \begin{cases} h/l(u) & \text{if } 1 - d/4 < s < 1 \\ (h/l(u)) |\ln(l(u)/h)|^{1/(4-4s)} & \text{if } 0 < s = 1 - d/4 \\ (h/l(u))^{d/(4-4s)} & \text{if } 0 < s < 1 - d/4 \end{cases}$$

and note that $\delta_u \leq 1/2$ if $h \leq C^{-1} l_0$ by (17). Moreover, (51) is an easy consequence of the corresponding estimate for $l(u)/l(u')$. With this choice we arrive at the bounds

$$I_u \leq C \begin{cases} h^{-d+2} l(u)^{-2} & \text{if } 1 - d/4 < s < 1 \\ h^{-d+2} l(u)^{-2} |\ln(l(u)/h)|^{1/2} & \text{if } 0 < s = 1 - d/4 \\ h^{-d/2+2s} l(u)^{-d/2-2s} & \text{if } 0 < s < 1 - d/4 \end{cases}.$$

Finally, we integrate with respect to u . The same arguments that lead to (18) and (19) yield

$$\int_{\Omega^*} I_u du \leq C \begin{cases} h^{-d+2} l_0^{-1} & \text{if } 1 - d/4 < s < 1 \\ h^{-d+2} l_0^{-1} |\ln(l_0/h)|^{1/2} & \text{if } 0 < s = 1 - d/4 \\ h^{-d/2+2s} l_0^{-d/2-2s+1} & \text{if } 0 < s < 1 - d/4 \end{cases}.$$

This completes the proof of the lower bound with the remainder stated in Proposition 3.

To prove the upper bound we put

$$\gamma = \int_{\mathbb{R}^d} \phi_u (\phi_u H_\Omega \phi_u)_-^0 \phi_u l(u)^{-d} du.$$

Obviously, $\gamma \geq 0$ holds and in view of (13) also $\gamma \leq 1$. The range of γ belongs to $\mathcal{H}^s(\Omega)$ and by the variational principle it follows that

$$-\text{Tr}(H_\Omega)_- \leq \text{Tr} \gamma H_\Omega = - \int_{\mathbb{R}^d} \text{Tr} (\phi_u H_\Omega \phi_u)_- l(u)^{-d} du.$$

This yields the upper bound and finishes the proof of Proposition 3. \square

6. DISCUSSION OF THE SECOND TERM

6.1. Representations for the second constant. In this section we study the second term of (6) in more detail. First we derive representation (8).

Proposition 17. *One has*

$$\begin{aligned} L_{s,d}^{(2)} &= \int_{\mathbb{R}^{d-1}} \zeta(|p'|^{-2s}) \frac{dp'}{(2\pi)^{d-1}} \\ &= \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2s}{(d-1)(d-1+2s)} \operatorname{Tr} [\chi A^{-(d-1)/2s} \chi - (A^+)^{-(d-1)/2s}]. \end{aligned} \quad (55)$$

Here χ is the characteristic function of \mathbb{R}_+ and

$$\zeta(\mu) = \mu^{-1} \int_0^\infty (a(\mu) - a^+(t, \mu)) dt. \quad (56)$$

Proof. The first identity follows immediately from (27) and (28). The second identity follows from the fact that

$$\int_{\mathbb{R}^{d-1}} |p'|^{2s} (E - |p'|^{-2s})_- \frac{dp'}{(2\pi)^{d-1}} = \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2s}{(d-1)(d-1+2s)} E^{-(d-1)/2s}$$

for any $E > 0$, which by the spectral theorem implies that

$$\int_{\mathbb{R}^{d-1}} |p'|^{2s} a^+(t, |p'|^{-2s}) \frac{dp'}{(2\pi)^{d-1}} = \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2s}{(d-1)(d-1+2s)} (A^+)^{-(d-1)/2s}(t, t)$$

and similarly for A . \square

Remark. There is another representation, namely,

$$L_{s,d}^{(2)} = \frac{2s}{d-1+2s} \int_{\mathbb{R}^{d-1}} \xi(|p'|^{-2s}) \frac{dp'}{(2\pi)^{d-1}}, \quad (57)$$

where

$$\xi(\mu) = \int_0^\infty (e(\mu) - e^+(t, \mu)) dt. \quad (58)$$

Here $e(\mu)$ and $e^+(t, \mu)$ are the diagonals of the integral kernels of the spectral projectors $(A - \mu)_-^0$ and $(A^+ - \mu)_-^0$, respectively. We have not shown that the integral in (58) converges, since we will not use (57) in the remainder of this paper. Identity (57) is an easy consequence of (55) and the fact that

$$a(\mu) = \int_0^\mu e(\tau) d\tau \quad a^+(t, \mu) = \int_0^\mu e^+(t, \tau) d\tau$$

which follows by the spectral theorem from $(E - \mu)_- = \int_0^\mu (E - \tau)_-^0 d\tau$. Representation (57) is natural since in terms of this function the conjectured formula for the *number* of

negative eigenvalues of H_Ω takes the form

$$\iint_{T\Omega} (|p|^{2s} - 1)_- \frac{dpdx}{(2\pi h)^d} - \iint_{T\partial\Omega} \xi(|p'|^{-2s}) \frac{dp'd\sigma(x)}{(2\pi h)^{d-1}} + o(h^{-d+1}),$$

which is the analogue of well-known two-term semi-classical formulas in the local case; see, for instance, [Ivr80, SV97]. The function ξ plays the role of a spectral shift. Note that we avoided to write (56) and (58) in terms of a trace. While the integrals on the diagonals converge, we do not expect the operators to be trace class, see [Pus08].

Remark. Yet another representation is

$$h^{-d+1} L_{s,d}^{(2)} = \int_0^\infty (H_-(x, x) - (H^+)_-(x, x)) dx_d.$$

(Note that the right side is independent of x' .) This follows from (30) and the corresponding formula for H . Using this representation one sees that our asymptotic formula coincides with the one obtained in [BK08, BKS09].

Finally, we refer to (71) in Section 8 for a representation of $L_{s,d}^{(2)}$ in terms of generalized eigenfunctions of A^+ .

6.2. Positivity of the constant. Here we shall prove

Proposition 18. *For any $0 < s < 1$ and $d \geq 2$, one has $L_{s,d}^{(2)} > 0$.*

Proof. We use the second representation in (55) and the fact that

$$E^{-(d-1)/2s} = \frac{1}{\Gamma((d-1)/2s)} \int_0^\infty e^{-\beta E} \beta^{(d-1)/(2s)-1} d\beta$$

for every $E > 0$ to see that

$$\begin{aligned} L_{s,d}^{(2)} &= \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2s}{(d-1)(d-1+2s)\Gamma((d-1)/2s)} \\ &\quad \times \int_0^\infty \int_0^\infty (e^{-\beta A}(t, t) - e^{-\beta A^+}(t, t)) \beta^{(d-1)/(2s)-1} dt d\beta. \end{aligned}$$

By means of the Trotter's product formula (see also [BK08] for a probabilistic derivation) it is easy to see that

$$\exp(-\beta A)(t, u) \geq \exp(-\beta A^+)(t, u)$$

for every $\beta > 0$ and every $t, u > 0$. This, together with the fact that $A \not\equiv A^+$, proves the proposition. \square

6.3. Comparison with a fractional power of the Dirichlet Laplacian. It is well-known that the Dirichlet Laplacian $-\Delta_\Omega$ on Ω satisfies

$$\mathrm{Tr} \left(-h^2 \Delta_\Omega - 1 \right)_- = L_{1,d}^{(1)} |\Omega| h^{-d} - L_{1,d}^{(2)} |\partial\Omega| h^{-d+1} + o(h^{-d+1}),$$

see, e.g., [FG11] for a proof under the sole assumption that $\partial\Omega \in C^{1,\alpha}$ for some $0 < \alpha \leq 1$. Here

$$L_{1,d}^{(1)} = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} (|p|^2 - 1)_- dp$$

and, by an argument similar to that in our Proposition 17, one can bring the second constant in the form

$$L_{1,d}^{(2)} = \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2}{(d-1)(d+1)} \mathrm{Tr} \left[\chi B^{-(d-1)/2} \chi - (B^+)^{-(d-1)/2} \right]$$

where $B = -\frac{d^2}{dt^2} + 1$ in $L^2(\mathbb{R})$ and $B^+ = -\frac{d^2}{dt^2} + 1$ with Dirichlet boundary conditions in $L^2(\mathbb{R}_+)$. A short computation, using the fact that

$$(E^s - 1)_- = s(1-s) \int_0^1 (E - \tau)_- \tau^{s-2} d\tau + s(E - 1)_-,$$

gives

$$\begin{aligned} \mathrm{Tr} \left((-h^2 \Delta_\Omega)^s - 1 \right)_- &= L_{1,d}^{(1)} |\Omega| h^{-d} s \left((1-s) \int_0^1 \tau^{d/2+s-1} d\tau + 1 \right) \\ &\quad - L_{1,d}^{(2)} |\partial\Omega| h^{-d+1} s \left((1-s) \int_0^1 \tau^{(d-1)/2+s-1} d\tau + 1 \right) \\ &\quad + o(h^{-d+1}) \\ &= L_{s,d}^{(1)} |\Omega| h^{-d} - \frac{s(d+1)}{d-1+2s} L_{1,d}^{(2)} |\partial\Omega| h^{-d+1} + o(h^{-d+1}), \end{aligned}$$

that is,

$$\tilde{L}_{s,d}^{(2)} = \frac{s(d+1)}{d-1+2s} L_{1,d}^{(2)} = \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2s}{(d-1)(d-1+2s)} \mathrm{Tr} \left[\chi B^{-(d-1)/2} \chi - (B^+)^{-(d-1)/2} \right].$$

Since

$$B^{-(d-1)/2}(t, t) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{(1+p^2)^{(d-1)/2}} dp = A^{-(d-1)/2s}(t, t)$$

we find that

$$\tilde{L}_{s,d}^{(2)} - L_{s,d}^{(2)} = \frac{|\mathbb{S}^{d-2}|}{(2\pi)^{d-1}} \frac{2s}{(d-1)(d-1+2s)} \mathrm{Tr} \left[(A^+)^{-(d-1)/2s} - (B^+)^{-(d-1)/2} \right].$$

We now apply Lemma 19 below with $B = -d^2/dt^2 + 1$ in $L^2(\mathbb{R})$, with P being the projector onto $L^2(\mathbb{R}_+)$ and with $\phi(E) = E^s$. Then $\phi(PBP) = (B^+)^s$ and $P\phi(B)P = A^+$, and therefore (59) yields

$$(B^+)^s \geq A^+.$$

Since $E \mapsto E^{-(d-1)/2s}$ is strictly monotone and since the operators A^+ and $(B^+)^s$ are not identical, we conclude that

$$\mathrm{Tr} \left((A^+)^{-(d-1)/2s} - (B^+)^{-(d-1)/2} \right) > 0.$$

This shows that $\tilde{L}_{s,d}^{(2)} - L_{s,d}^{(2)} > 0$ and completes the proof of Proposition 2. \square

In the previous proof we used

Lemma 19. *Let B be a non-negative operator with $\ker B = \{0\}$ and let P be an orthogonal projection. Then for any complete Bernstein function*

$$\phi(PBP) \geq P\phi(B)P. \quad (59)$$

We recall (see, e.g., [SSV10]) that complete Bernstein functions (also known as operator-monotone functions) are characterized by the representation

$$\phi(E) = a + bE + \int_0^\infty \frac{E}{\tau(E + \tau)} d\rho(\tau) \quad (60)$$

with $a \in \mathbb{R}$, $b \geq 0$ and a positive measure ρ satisfying

$$\int_0^\infty \frac{1}{\tau(1 + \tau)} d\rho(\tau) < \infty.$$

Proof. We first prove that

$$B^{-1} \geq (PBP)^{-1}. \quad (61)$$

Let us write $P^\perp = 1 - P$, so that

$$B = PBP + P^\perp BP^\perp + P^\perp BP + PBP^\perp.$$

By the Schwarz inequality we have

$$P^\perp BP + PBP^\perp \leq \varepsilon PBP + \frac{1}{\varepsilon} P^\perp BP^\perp$$

for any $\varepsilon > 0$, and hence

$$B \leq (1 + \varepsilon)PBP + \left(1 + \frac{1}{\varepsilon}\right) P^\perp BP^\perp.$$

Since P and P^\perp are orthogonal we can invert this inequality and obtain

$$B^{-1} \geq \frac{1}{1 + \varepsilon} (PBP)^{-1} + \frac{\varepsilon}{1 + \varepsilon} (P^\perp BP^\perp)^{-1}.$$

Thus (61) follows by taking $\varepsilon \rightarrow 0+$.

Now if ϕ is of the form (60) then

$$\phi(PBP) - P\phi(B)P = - \int_0^\infty \left((PBP + \tau)^{-1} - P(B + \tau)^{-1}P \right) d\rho(\tau).$$

By (61) with B replaced by $B + \tau$, this is non-negative. \square

7. EQUIVALENCE OF (2) AND (3)

For the sake of completeness we include a short proof of

Lemma 20. *Let $(\lambda_k)_{k \in \mathbb{N}}$ be a non-decreasing sequence of real numbers and let $A, C > 0$, $B, D \in \mathbb{R}$ and $-1 < a - 1 < b < a$ be related by*

$$C = A^{-1/a} a(a+1)^{-(1+a)/a}, \quad D = B(A(a+1))^{-(1+b)/a}.$$

Then the asymptotic formula

$$\sum_{k=1}^N \lambda_k = AN^{a+1} + BN^{b+1}(1 + o(1)), \quad N \rightarrow \infty, \quad (62)$$

is equivalent to

$$\sum_{k \in \mathbb{N}} (\Lambda - \lambda_k)_+ = C\Lambda^{(1+a)/a} - D\Lambda^{(1+b)/a}(1 + o(1)), \quad \Lambda \rightarrow \infty. \quad (63)$$

Proof. This lemma is a consequence of Hardy, Littlewood and Polya's majorization theorem, which says that for any non-decreasing sequences $\{a_k\}$ and $\{b_k\}$

$$\sum_{k=1}^N a_k \leq \sum_{k=1}^N b_k \quad \text{for all } N \in \mathbb{N} \quad (64)$$

is equivalent to

$$\sum_{k=1}^{\infty} (\Lambda - a_k)_+ \geq \sum_{k=1}^{\infty} (\Lambda - b_k)_+ \quad \text{for all } \Lambda \in \mathbb{R};$$

see, e.g., [MO79, Prop. 4.B.4]. As usual, we will denote property (64) by $\{a_k\} \prec \{b_k\}$.

We fix $\epsilon > 0$ and set $\beta_k^\pm = A(a+1)k^a + (B \pm \epsilon)(b+1)k^b$. Note that the assumptions on a and b imply

$$\sum_{k=1}^N \beta_k^\pm = AN^{a+1} + (B \pm \epsilon)N^{b+1}(1 + o(1)), \quad N \rightarrow \infty, \quad (65)$$

and

$$\sum_{k \in \mathbb{N}} (\Lambda - \beta_k^\pm)_+ = \frac{aA}{(A(a+1))^{1+1/a}} \Lambda^{(1+a)/a} - \frac{B \pm \epsilon}{(A(a+1))^{(1+b)/a}} \Lambda^{(1+b)/a}(1 + o(1)), \quad (66)$$

as $\Lambda \rightarrow \infty$. First, we assume that (62) holds. Then, by (62) and (65) there is an $N_\epsilon \in \mathbb{N}$ such that for all $N \geq N_\epsilon$

$$\sum_{k=1}^N \beta_k^- \leq \sum_{k=1}^N \lambda_k \leq \sum_{k=1}^N \beta_k^+.$$

We put $\alpha_k^\pm = \beta_k^\pm$ for $k \geq N_\epsilon$ and $\alpha_k^+ = \max(\beta_k^+, \lambda_k)$, $\alpha_k^- = \min(\beta_k^-, \lambda_k)$ for $k < N_\epsilon$. Thus $\{\alpha_k^-\} \prec \{\lambda_k\} \prec \{\alpha_k^+\}$, and therefore

$$\sum_{k \in \mathbb{N}} (\Lambda - \alpha_k^+)_+ \leq \sum_{k \in \mathbb{N}} (\Lambda - \lambda_k)_+ \leq \sum_{k \in \mathbb{N}} (\Lambda - \alpha_k^-)_+ \quad \text{for all } \Lambda \in \mathbb{R}.$$

Since $\sum_{k \in \mathbb{N}} (\Lambda - \alpha_k^\pm)_+ = \sum_{k \in \mathbb{N}} (\Lambda - \beta_k^\pm)_+ + O(1)$, the assertion (63) follows from (66). The converse implication is proved similarly. \square

8. THE ONE-DIMENSIONAL MODEL OPERATOR

Here we outline the calculations that are necessary to complete the analysis of the model operator A^+ introduced in Section 3. The results depend on the following spectral representation of the operator A^+ found in [Kwa10b].

Theorem 21. *For $E > 0$ let*

$$\psi(E) = (E + 1)^s - 1$$

and for $\lambda > 0$ put $\gamma_\lambda(\xi) = 0$ if $0 < \xi < 1$ and

$$\begin{aligned} \gamma_\lambda(\xi) &= \frac{1}{\pi} \frac{\lambda \psi'(\lambda^2) \sin(\pi s) (\xi^2 - 1)^s}{\psi(\lambda^2)^2 + (\xi^2 - 1)^s - 2\psi(\lambda^2)(\xi^2 - 1) \cos(\pi s)} \\ &\quad \times \exp\left(-\frac{1}{\pi} \int_0^\infty \frac{\xi}{\xi^2 + \zeta^2} \ln \frac{\psi'(\lambda^2)(\lambda^2 - \zeta^2)}{\psi(\lambda^2) - \psi(\zeta^2)} d\zeta\right) \end{aligned}$$

if $\xi \geq 1$. Moreover, define a phase-shift

$$\vartheta_\lambda = \frac{1}{\pi} \int_0^\infty \frac{\lambda}{\zeta^2 - \lambda^2} \ln \frac{\psi'(\lambda^2)(\lambda^2 - \zeta^2)}{\psi(\lambda^2) - \psi(\zeta^2)} d\zeta \quad (67)$$

and functions

$$F_\lambda(x) = \sin(\lambda x + \vartheta_\lambda) + \int_0^\infty e^{-x\xi} \gamma_\lambda(\xi) d\xi, \quad x > 0. \quad (68)$$

Then

$$\Phi f(\lambda) = \sqrt{\frac{2}{\pi}} \int_0^\infty f(x) F_\lambda(x) dx$$

defines a unitary operator from $L^2(\mathbb{R}_+)$ to $L^2(\mathbb{R}_+)$.

This operator diagonalizes A^+ in the sense that a function $f \in L^2(\mathbb{R}_+)$ is in the domain of A^+ if and only if $(\lambda^2 + 1)^s \Phi f(\lambda)$ is in $L^2(\mathbb{R}_+)$, and in this case

$$\Phi A^+ f(\lambda) = (\lambda^2 + 1)^s \Phi f(\lambda).$$

According to [Kwa10b] the Laplace transform of γ_λ is a completely monotone function bounded by one. From (68) it follows that for all $t \geq 0$

$$|F_\lambda(t)| \leq 2. \quad (69)$$

Theorem 21 states that the functions F_λ are generalized eigenfunctions of the operator A^+ . Hence, we can write

$$e^+(t, u, \mu) = \frac{2}{\pi} \int_0^\infty ((\lambda^2 + 1)^s - \mu)_-^0 F_\lambda(t) F_\lambda(u) d\lambda. \quad (70)$$

From (28) and (8) it follows that

$$L_{s,d}^{(2)} = \frac{4s}{(d-1+2s)(d-1)} \frac{|\mathbb{S}^{d-2}|}{(2\pi)^d} \int_0^\infty \int_0^\infty (1 - 2F_\lambda^2(t)) (\lambda^2 + 1)^{-(d-1)/2} d\lambda dt. \quad (71)$$

8.1. **Proof of Lemma 9.** Lemma 9 is an immediate consequence of (70). In view of (69) we estimate

$$|e^+(t, u, \mu)| \leq C \int_0^{(\mu^{1/s}-1)_+^{1/2}} d\lambda \leq C\mu^{1/(2s)}.$$

This proves the lemma.

8.2. **Proof of Lemma 10.** First we need the following technical result about ϑ_λ .

Lemma 22. *The phase-shift ϑ_λ is monotone increasing and twice differentiable in $\lambda > 0$. It satisfies*

$$\vartheta_0 = 0 \quad \text{and} \quad \vartheta_\lambda \rightarrow \frac{\pi}{4}(1-s) \quad \text{as} \quad \lambda \rightarrow \infty.$$

The first and second derivatives are bounded and one has, as $\lambda \rightarrow \infty$,

$$\frac{d\vartheta_\lambda}{d\lambda} = \frac{d^2\vartheta_\lambda}{d\lambda^2} = O\left(\frac{1}{\lambda}\right).$$

Proof. Following [Kwa10b], we substitute $\zeta = \lambda z$ for $\zeta \in (0, 1)$ and $\zeta = \lambda/z$ for $\zeta \in (1, \infty)$ in the definition of ϑ_λ and obtain

$$\vartheta_\lambda = \frac{1}{\pi} \int_0^1 \frac{1}{1-z^2} \ln \left(\frac{1}{z^2} \frac{\psi(\lambda^2) - \psi(\lambda^2 z^2)}{\psi(\lambda^2/z^2) - \psi(\lambda^2)} \right) dz.$$

Note that the function

$$\frac{1}{z^2} \frac{\psi(\lambda^2) - \psi(\lambda^2 z^2)}{\psi(\lambda^2/z^2) - \psi(\lambda^2)} = \frac{1}{z^2} \frac{(1+\lambda^2)^s - (1+\lambda^2 z^2)^s}{(1+\lambda^2/z^2)^s - (1+\lambda^2)^s}$$

equals 1 for $\lambda = 0$ and that for all $z \in (0, 1)$ it is increasing in $\lambda > 0$ and tends to z^{2s-2} as λ tends to infinity. By Lebesgue's dominated convergence we find $\vartheta_0 = 0$ and

$$\lim_{\lambda \rightarrow \infty} \vartheta_\lambda = \frac{1}{\pi} \int_0^1 \frac{1}{1-z^2} \ln(z^{2s-2}) dz = \frac{\pi}{4}(1-s).$$

By (67), we also have

$$\vartheta_\lambda = \frac{1}{\pi} \int_0^\infty b_\lambda(\zeta) d\zeta$$

with

$$b_\lambda(\zeta) = \frac{\lambda}{\zeta^2 - \lambda^2} \ln \left(\frac{s(1+\lambda^2)^{s-1}(\lambda^2 - \zeta^2)}{(\lambda^2 + 1)^s - (\zeta^2 + 1)^s} \right).$$

We remark that

$$|\partial_\lambda b_\lambda(\zeta)| \leq \partial_\lambda b_\lambda(\zeta)|_{\lambda=0} = \frac{1}{\zeta^2} \ln \left(\frac{s\zeta^2}{(1-\zeta^2)^s - 1} \right).$$

for all $\zeta \in (0, \infty)$. Since the last expression is integrable in $\zeta \in (0, \infty)$ it follows that

$$\frac{d\vartheta_\lambda}{d\lambda} = \frac{1}{\pi} \int_0^\infty \partial_\lambda b_\lambda(\zeta) d\lambda$$

is bounded and, in particular, we obtain

$$\left. \frac{d\vartheta_\lambda}{d\lambda} \right|_{\lambda=0} = \frac{1}{\pi} \int_0^\infty \frac{1}{\zeta^2} \ln \left(\frac{s\zeta^2}{(1+\zeta^2)^s - 1} \right) d\zeta. \quad (72)$$

Similarly, we can show existence and boundedness of the second derivative and decay of the derivatives as $\lambda \rightarrow \infty$ by explicit calculations and Lebesgue's dominated convergence. \square

To simplify notation we put

$$\psi_\lambda(E) = \frac{1 - E/\lambda^2}{1 - \psi(E)/\psi(\lambda^2)}$$

for $E > 0$. Moreover, we write G_λ for the Laplace transform of γ_λ and g_λ for the Laplace transform of G_λ . According to [Kwa10b] we have

$$g_\lambda(t) = \frac{\lambda \cos \vartheta_\lambda + t \sin \vartheta_\lambda}{\lambda^2 + t^2} - \lambda^2 \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)} \frac{\varphi_\lambda(t)}{\lambda^2 + t^2}}, \quad t > 0, \quad (73)$$

with

$$\varphi_\lambda(t) = \exp \left(\frac{1}{\pi} \int_0^\infty \frac{t}{t^2 + \zeta^2} \ln (\psi_\lambda(\zeta^2)) d\zeta \right).$$

To prove Lemma 10 we need the following properties of φ_λ .

Lemma 23. *The function $t \mapsto \varphi_\lambda(t)$ is differentiable in $t > 0$ and its derivative satisfies*

$$\begin{aligned} \varphi'_\lambda(0) &= o(1) \quad \text{as } \lambda \rightarrow \infty, \\ \varphi'_\lambda(0) &= \left. \frac{d\vartheta_\lambda}{d\lambda} \right|_{\lambda=0} + O(\lambda) \quad \text{as } \lambda \rightarrow 0. \end{aligned}$$

Proof. For fixed $\zeta \in (0, \infty)$ the function $\lambda \mapsto \psi_\lambda(\zeta^2)$ is non-increasing in $\lambda > 0$ and tends to 1 as $\lambda \rightarrow \infty$. Moreover,

$$\frac{1}{\zeta^2} \ln (\psi_0(\zeta^2)) = \frac{1}{\zeta^2} \ln \left(\frac{s\zeta^2}{(\zeta^2 + 1)^s - 1} \right)$$

is integrable with respect to $\zeta \in (0, \infty)$. Hence we find that

$$\varphi'_\lambda(0) = \frac{1}{\pi} \int_0^\infty \frac{1}{\zeta^2} \ln (\psi_\lambda(\zeta^2)) d\zeta$$

and $\varphi'_\lambda(0) = o(1)$ as $\lambda \rightarrow \infty$ by Lebesgue's theorem.

In view of (72)

$$\varphi'_\lambda(0)|_{\lambda=0} = \frac{1}{\pi} \int_0^\infty \frac{1}{\zeta^2} \ln (\psi_0(\zeta^2)) d\zeta = \left. \frac{d\vartheta_\lambda}{d\lambda} \right|_{\lambda=0}.$$

The second claim now follows from the fact that the derivative of $\lambda \mapsto \varphi'_\lambda(0)$ is bounded. \square

Proof of Lemma 10. In view of Theorem 21 we can write

$$a(\mu) - a^+(t, \mu) = \frac{1}{\pi} \int_0^\infty ((\lambda^2 + 1)^s - \mu)_- (1 - 2F_\lambda^2(t)) d\lambda$$

and by (68)

$$1 - 2F_\lambda(t)^2 = \cos(2\lambda t + 2\vartheta_\lambda) - 4 \sin(\lambda t + \vartheta_\lambda) G_\lambda(t) - 2G_\lambda(t)^2.$$

We get

$$\int_0^\infty t^\gamma |a(\mu) - a^+(t, \mu)| dt \leq R_1(\mu) + R_2(\mu)$$

with

$$R_1(\mu) = \int_0^\infty t^\gamma \left| \int_0^{(\mu^{1/s} - 1)_+^{1/2}} (\mu - (\lambda^2 + 1)^s) \cos(2\lambda t + 2\vartheta_\lambda) d\lambda \right| dt,$$

$$R_2(\mu) = \int_0^\infty t^\gamma \left| \int_0^{(\mu^{1/s} - 1)_+^{1/2}} (\mu - (\lambda^2 + 1)^s) (2 \sin(\lambda t + \vartheta_\lambda) G_\lambda(t) + G_\lambda(t)^2) d\lambda \right| dt.$$

To estimate $R_1(\mu)$ we split the integration in t and integrate over $t \in [0, 1]$ first. We assume $0 < \gamma < 1$. The proof for $\gamma = 0$ follows similarly.

We write

$$\cos(2\lambda t + 2\vartheta_\lambda) = \frac{1}{2t} \frac{d}{d\lambda} \sin(2\lambda t + 2\vartheta_\lambda) - \frac{\cos(2\lambda t + 2\vartheta_\lambda)}{t} \frac{d\vartheta_\lambda}{d\lambda}$$

and insert this identity in the expression for $R_1(\mu)$. After integrating by parts in the λ -integral one can estimate

$$\int_0^1 t^\gamma \left| \int_0^{(\mu^{1/s} - 1)_+^{1/2}} (\mu - (\lambda^2 + 1)^s) \cos(2\lambda t + 2\vartheta_\lambda) d\lambda \right| dt \leq C\mu ((\ln \mu)^2 + 1).$$

To estimate the integral over $t \in [1, \infty)$ we proceed similarly. We integrate by parts twice and get

$$\int_1^\infty t^\gamma \left| \int_0^{(\mu^{1/s} - 1)_+^{1/2}} (\mu - (\lambda^2 + 1)^s) \cos(2\lambda t + 2\vartheta_\lambda) d\lambda \right| dt \leq C\mu (\ln \mu + 1).$$

We conclude

$$R_1(\mu) \leq C\mu ((\ln \mu)^2 + 1)$$

and turn to estimating $R_2(\mu)$.

Since G_λ is non-negative and uniformly bounded, we have

$$R_2(\mu) \leq C \int_0^{(\mu^{1/s} - 1)_+^{1/2}} (\mu - (\lambda^2 + 1)^s) \int_0^\infty t^\gamma G_\lambda(t) du d\lambda. \quad (74)$$

Identity (73) implies $\int_0^\infty G_\lambda(t)dt = g_\lambda(0)$ and $\int_0^\infty tG_\lambda(t)dt = g'_\lambda(0)$. We note that

$$g_\lambda(0) = \frac{\cos \vartheta_\lambda}{\lambda} - \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}$$

and apply Lemma 22 to estimate $\int_0^\infty G_\lambda(t)dt \leq C(\lambda \wedge \lambda^{-1})$. Moreover, by (73),

$$g'_\lambda(0) = \frac{\sin \vartheta_\lambda}{\lambda^2} - \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}} \varphi'_\lambda(0)$$

and we apply Lemma 22 and Lemma 23 to estimate $\int_0^\infty tG_\lambda(t)dt \leq C(1 \wedge \lambda^{-1})$. It follows that

$$\int_0^\infty t^\gamma G_\lambda(t) dt \leq C(1 \wedge \lambda^{-1}).$$

Thus, by (74), we arrive at

$$R_2(\mu) \leq C \int_0^{(\mu^{1/s}-1)_+^{1/2}} (\mu - (\lambda^2 + 1)^s) (1 \wedge \lambda^{-1}) d\lambda \leq C\mu(\ln \mu + 1).$$

This finishes the first part of the proof of Lemma 10.

In order to prove the assertion about $K(t)$, we bound

$$\int_0^\infty t^\gamma |K(t)| dt \leq \int_{|\xi'| < 1} |\xi'|^{1+2s} \int_0^\infty t^\gamma |a^+(t|\xi'|, |\xi'|^{-2s}) - a(|\xi'|^{-2s})| dt d\xi'.$$

Here we also used that, since $a(\mu) = a^+(t, \mu) = 0$ for $\mu \leq 1$, we can restrict the integration in the definition of K to $|\xi'| < 1$. On the other hand, from (26) we know that

$$\int_0^\infty t^\gamma |a^+(t\mu^{-1/2s}, \mu) - a(\mu)| dt \leq C_\gamma \mu^{1+(\gamma+1)/(2s)} ((\ln \mu)^2 + 1).$$

Combining these two bounds and using that $\gamma < 1 \leq d-1$ we obtain the second part of Lemma 10. \square

8.3. A remainder estimate. The following technical lemma was needed in the proof of the upper bound near the boundary.

Lemma 24. *Assume that $\phi \in C_0^1(\mathbb{R}^d)$ is supported in a ball of radius $l = 1$ and that (14) is satisfied with $l = 1$. Then for any $\frac{1}{2} - s < \sigma < \min\{\frac{1}{2}, 1 - s\}$ one has*

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \left| (-\Delta_{x'})^\sigma \frac{|\phi(x) - \phi(y)|^2}{|x - y|^{d+2s}} \right| dx dy \leq C \quad (75)$$

Proof. For $x = (x', x_d) \in \mathbb{R}^{d-1} \times \mathbb{R}$ and $y = (y', y_d) \in \mathbb{R}^{d-1} \times \mathbb{R}$ put

$$F_{x_d, y}(x') = \frac{(\phi(x', x_d) - \phi(y', y_d))^2}{(|x' - y'|^2 + (x_d - y_d)^2)^{d/2+s}}.$$

To establish (75) we use that

$$\left| (-\Delta_{x'})^\sigma \frac{|\phi(x) - \phi(y)|^2}{|x - y|^{d+2s}} \right| \leq C \int_{\mathbb{R}^{d-1}} \frac{|F_{x_d,y}(x') - F_{x_d,y}(z')|}{|x' - z'|^{d-1+2\sigma}} dz' \quad (76)$$

and split the integration in $x \in \mathbb{R}^d$ and $y \in \mathbb{R}^d$ in four parts. First we assume that x and y are in B_1 . Then we have to show that

$$\begin{aligned} & \int_{B_1} \int_{B_1} \int_{\mathbb{R}^{d-1}} \frac{|F_{x_d,y}(x') - F_{x_d,y}(z')|}{|x' - z'|^{d-1+2\sigma}} dz' dx dy = \\ & \int_{B_1} \int_{B_1} \int_{|x' - z'| < |x - y|/2} \frac{|F_{x_d,y}(x') - F_{x_d,y}(z')|}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \\ & + \int_{B_1} \int_{B_1} \int_{|x' - z'| \geq |x - y|/2} \frac{|F_{x_d,y}(x') - F_{x_d,y}(z')|}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \end{aligned} \quad (77)$$

is bounded from above.

To estimate the first integral over $|x' - z'| < |x - y|/2$ we use that

$$F(z') - F(x') = \sum_{j=1}^{d-1} \frac{(z_j - x_j)}{|x' - z'|} \int_0^{|x' - z'|} (\partial_j F) \left(x' + t \frac{(z' - x')}{|x' - z'|} \right) dt.$$

For $j = 1, \dots, d-1$ we have

$$(\partial_j F_{x_d,y})(x') = \frac{2(\phi(x', x_d) - \phi(y))(\partial_j \phi(x))}{|x - y|^{d+2s}} - (d+2s)(x_j - y_j) \frac{(\phi(x) - \phi(y))^2}{|x - y|^{d+2s+2}},$$

thus

$$|(\partial_j F_{x_d,y})(x')| \leq C |x - y|^{-d+1-2s}.$$

Hence, we obtain

$$\begin{aligned} & |F_{x_d,y}(z') - F_{x_d,y}(x')| \\ & \leq C |x' - z'|^\alpha \left(\int_0^{|x' - z'|} \left(\left| x' + t \frac{(z' - x')}{|x' - z'|} - y' \right|^2 + (x_d - y_d)^2 \right)^\beta dt \right)^{1-\alpha}, \end{aligned} \quad (78)$$

with $0 < \alpha < 1$ and $\beta = (\frac{d-1}{2} + s)/(\alpha - 1)$, by applying Hölder's inequality. Note that

$$\begin{aligned} \left| x' - y' + t \frac{(z' - x')}{|x' - z'|} \right|^2 + (x_d - y_d)^2 &= |x - y|^2 + t^2 + 2t \frac{(x' - y') \cdot (z' - x')}{|x' - z'|} \\ &\geq (|x - y| - t)^2. \end{aligned}$$

Inserting this into (78) we get for $|x' - z'| < |x - y|/2$

$$\begin{aligned} |F_{x_d,y}(z') - F_{x_d,y}(x')| &\leq C |x' - z'|^\alpha \left(\int_0^{|x-y|/2} (|x - y| - t)^{2\beta} dt \right)^{1-\alpha} \\ &\leq C |x' - z'|^\alpha |x - y|^{(2\beta+1)(1-\alpha)}, \end{aligned}$$

where $(2\beta + 1)(1 - \alpha) = -d - 2s + 2 - \alpha$. We conclude that for any $2\sigma < \alpha < 1$ and $\sigma < 1 - s$

$$\begin{aligned} & \int_{B_1} \int_{B_1} \int_{|x'-z'| < |x-y|/2} \frac{|F_{x_d,y}(x') - F_{x_d,y}(z')|}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \\ & \leq C \int_{B_1} \int_{B_1} \int_{|x'-z'| < |x-y|/2} |x' - z'|^{-d+1-2\sigma+\alpha} dz' |x - y|^{-d-2s+2-\alpha} dx dy \\ & \leq C. \end{aligned} \quad (79)$$

Now we turn to the second integral in (77) over $|x' - z'| \geq |x - y|/2$. Since

$$0 \leq F_{x_d,y}(x') \leq |x - y|^{-d-2s+2} \quad (80)$$

and $\sigma < 1 - s$ we have

$$\int_{B_1} \int_{B_1} \int_{|x'-z'| \geq |x-y|/2} \frac{F_{x_d,y}(x')}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \leq C \int_{B_1} \int_{B_1} \frac{1}{|x - y|^{d+2s-2+2\sigma}} \leq C. \quad (81)$$

Moreover,

$$\int_{|x'-z'| \geq |x-y|/2} \frac{F_{x_d,y}(z')}{|x' - z'|^{d-1+2\sigma}} dz' \leq C |x - y|^{1-d-2\sigma+(d-1)/p} \left(\int_{|x'-z'| \geq |x-y|/2} F_{x_d,y}^q(z') dz' \right)^{\frac{1}{q}}$$

with $\frac{1}{p} + \frac{1}{q} = 1$, by Hölder's inequality. Since $\sigma > \frac{1}{2} - s$ we can choose $p > \frac{d-1}{2\sigma}$ and $q > \frac{d-1}{d+2s-2}$. By (80), we have

$$\begin{aligned} \left(\int_{|x'-z'| \geq |x-y|/2} F_{x_d,y}^q(z') dz' \right)^{\frac{1}{q}} & \leq C \left(\int_{\mathbb{R}^{d-1}} (|z' - y_d|^2 + (x_d - y_d)^2)^{-q(d/2+s-1)} dz' \right)^{\frac{1}{q}} \\ & \leq C |x_d - y_d|^{-d-2s+2+(d-1)/q}. \end{aligned}$$

It follows that

$$\begin{aligned} & \int_{B_1} \int_{B_1} \int_{|x'-z'| \geq |x-y|/2} \frac{F_{x_d,y}(z')}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \\ & \leq C \int_{B_1} \int_{B_1} |x - y|^{-d+1-2\sigma+(d-1)/p} |x_d - y_d|^{-d-2s+2+(d-1)/q} dx dy \\ & \leq C \int_0^2 t^{-d-2s+2+(d-1)/q} \int_0^2 r^{d-2} (r^2 + t^2)^{-(d+1-2\sigma)/2+(d-1)/(2p)} dr dt, \end{aligned}$$

where we substituted $t = |x_d - y_d|$ and $r = |x' - y_d|$. Since $p > \frac{d-1}{2\sigma}$ and $\sigma < 1 - s$ we find

$$\int_{B_1} \int_{B_1} \int_{|x'-z'| \geq |x-y|/2} \frac{F_{x_d,y}(z')}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \leq C \int_0^2 t^{1-2s-2\sigma} dt \leq C. \quad (82)$$

The estimates (81) and (82) show that

$$\int_{B_1} \int_{B_1} \int_{|x'-z'| \geq |x-y|/2} \frac{|F_{x_d,y}(x') - F_{x_d,y}(z')|}{|x' - z'|^{d-1+2\sigma}} dz' dx dy \leq C \quad (83)$$

and from (76), (79), and (83) it follows that

$$\int_{B_1} \int_{B_1} \left| (-\Delta_{x'})^\sigma \frac{|\phi(x) - \phi(y)|^2}{|x - y|^{d+2s}} \right| dx dy \leq C.$$

The proof that the respective integrals over $B_1 \times (\mathbb{R}^d \setminus B_1)$, $(\mathbb{R}^d \setminus B_1) \times B_1$, and $(\mathbb{R}^d \setminus B_1) \times (\mathbb{R}^d \setminus B_1)$ are finite is similar but easier, since $\text{supp} \phi \subset B_1$ and we only have to handle one singularity at a time. \square

REFERENCES

- [Ber72] F. A. Berezin, *Covariant and contravariant symbols of operators*, Izv. Akad. Nauk SSSR Ser. Mat. **13** (1972), 1134–1167.
- [BG59] R. M. Blumenthal and R. K. Gettoor, *The asymptotic distribution of the eigenvalues for a class of Markov operators*, Pacific J. Math. **9** (1959), 399–408.
- [BK08] R. Bañuelos and T. Kulczycki, *Trace estimates for stable processes*, Probab. Theory Related Fields **142** (2008), 313–338.
- [BKS09] R. Bañuelos, T. Kulczycki, and B. Siudeja, *On the trace of symmetric stable processes on Lipschitz domains*, J. Funct. Anal. **257** (2009), no. 10, 3329–3352.
- [CS05] Z. Chen and R. Song, *Two-sided eigenvalue estimates for subordinate processes in domains*, J. Funct. Anal. **226** (2005), no. 1, 90–113.
- [FG11] R. L. Frank and L. Geisinger, *Two-term spectral asymptotics of the Dirichlet Laplacian on a bounded domain*, Mathematical Results in Quantum Physics: Proceedings of the Qmath11 Conference (Pavel Exner, ed.), World Scientific Publishing Company, 2011, pp. 138–147.
- [Hör85] L. Hörmander, *The analysis of linear partial differential operators, vol. 4*, Springer-Verlag, Berlin, 1985.
- [Ivr80] V. Ja. Ivrii, *The second term of the spectral asymptotics for the Laplace-Beltrami operator on manifolds with boundary*, Funktsional. Anal. i Prilozhen. **14** (1980), no. 2, 25–34.
- [Kat66] T. Kato, *Perturbation theory for linear operators*, Grundlehren Math. Wiss., vol. 132, Springer-Verlag, New York, 1966.
- [KKMS10] T. Kulczycki, M. Kwaśnicki, J. Małecki, and A. Stos, *Spectral properties of the Cauchy process on half-line and interval*, Proc. Lond. Math. Soc. (3) **101** (2010), no. 2, 589–622.
- [Kwa10a] M. Kwasnicki, *Eigenvalues of the fractional Laplace operator in the interval*, arXiv:1012.1133v1 (2010).
- [Kwa10b] ———, *Spectral analysis of subordinate Brownian motions in half-line*, arXiv:1006.0524v2 (2010).
- [Lie73] E. H. Lieb, *The classical limit of quantum spin systems*, Comm. Math. Phys. **31** (1973), 327–340.
- [LY83] P. Li and S. T. Yau, *On the Schrödinger equation and the eigenvalue problem*, Comm. Math. Phys. **88** (1983), no. 3, 309–318.
- [LY88] E. H. Lieb and H. T. Yau, *The stability and instability of relativistic matter*, Comm. Math. Phys. **118** (1988), no. 2, 177–213.
- [MO79] A. W. Marshall and I. Olkin, *Inequalities: theory of majorization and its applications*, Mathematics in Science and Engineering, vol. 143, Academic Press Inc., New York, 1979.
- [Pus08] A. Pushnitski, *The scattering matrix and the differences of spectral projections*, Bull. Lond. Math. Soc. **40** (2008), no. 2, 227–238.
- [SS03] J. P. Solovej and W. L. Spitzer, *A new coherent states approach to semiclassics which gives Scott’s correction*, Comm. Math. Phys. **241** (2003), 383–420.
- [SSS10] J. P. Solovej, T. Østergaard Sørensen, and W. L. Spitzer, *The relativistic Scott correction for atoms and molecules*, Comm. Pure Appl. Math. **63** (2010), 39–118.

- [SSV10] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein functions*, de Gruyter Studies in Mathematics, vol. 37, Walter de Gruyter & Co., Berlin, 2010, Theory and applications.
- [SV97] Y. Safarov and D. Vassiliev, *The asymptotic distribution of eigenvalues of partial differential operators*, Translations of Mathematical Monographs, 155, American Mathematical Society, Providence, RI, 1997.

Rupert L. Frank

Leander Geisinger
Universität Stuttgart
Pfaffenwaldring 57
D - 70569 Stuttgart
E-Mail: geisinger@mathematik.uni-stuttgart.de

Erschienenene Preprints ab Nummer 2007/001

Komplette Liste: <http://www.mathematik.uni-stuttgart.de/preprints>

- 2011/020 *Frank, R.L.; Geisinger, L.:* Refined Semiclassical Asymptotics for Fractional Powers of the Laplace Operator
- 2011/019 *Frank, R.L.; Geisinger, L.:* Two-term spectral asymptotics for the Dirichlet Laplacian on a bounded domain
- 2011/018 *Hänel, A.; Schulz, C.; Wirth, J.:* Embedded eigenvalues for the elastic strip with cracks
- 2011/017 *Wirth, J.:* Thermo-elasticity for anisotropic media in higher dimensions
- 2011/016 *Höllig, K.; Hörner, J.:* Programming Multigrid Methods with B-Splines
- 2011/015 *Ferrario, P.:* Nonparametric Local Averaging Estimation of the Local Variance Function
- 2011/014 *Müller, S.; Dippon, J.:* k-NN Kernel Estimate for Nonparametric Functional Regression in Time Series Analysis
- 2011/013 *Knarr, N.; Stroppel, M.:* Unitals over composition algebras
- 2011/012 *Knarr, N.; Stroppel, M.:* Baer involutions and polarities in Moufang planes of characteristic two
- 2011/011 *Knarr, N.; Stroppel, M.:* Polarities and planar collineations of Moufang planes
- 2011/010 *Jentsch, T.; Moroianu, A.; Semmelmann, U.:* Extrinsic hyperspheres in manifolds with special holonomy
- 2011/009 *Wirth, J.:* Asymptotic Behaviour of Solutions to Hyperbolic Partial Differential Equations
- 2011/008 *Stroppel, M.:* Orthogonal polar spaces and unitals
- 2011/007 *Nagl, M.:* Charakterisierung der Symmetrischen Gruppen durch ihre komplexe Gruppenalgebra
- 2011/006 *Solanes, G.; Teufel, E.:* Horo-tightness and total (absolute) curvatures in hyperbolic spaces
- 2011/005 *Ginoux, N.; Semmelmann, U.:* Imaginary Kählerian Killing spinors I
- 2011/004 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic D -Scales: Part II — Gain-Scheduled Control
- 2011/003 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic D -Scales: Part I — Robust Control
- 2011/002 *Alexandrov, B.; Semmelmann, U.:* Deformations of nearly parallel G_2 -structures
- 2011/001 *Geisinger, L.; Weidl, T.:* Sharp spectral estimates in domains of infinite volume
- 2010/018 *Kimmerle, W.; Konovalov, A.:* On integral-like units of modular group rings
- 2010/017 *Gauduchon, P.; Moroianu, A.; Semmelmann, U.:* Almost complex structures on quaternion-Kähler manifolds and inner symmetric spaces
- 2010/016 *Moroianu, A.; Semmelmann, U.:* Clifford structures on Riemannian manifolds
- 2010/015 *Grafarend, E.W.; Kühnel, W.:* A minimal atlas for the rotation group $SO(3)$
- 2010/014 *Weidl, T.:* Semiclassical Spectral Bounds and Beyond
- 2010/013 *Stroppel, M.:* Early explicit examples of non-desarguesian plane geometries
- 2010/012 *Effenberger, F.:* Stacked polytopes and tight triangulations of manifolds
- 2010/011 *Györfi, L.; Walk, H.:* Empirical portfolio selection strategies with proportional transaction costs
- 2010/010 *Kohler, M.; Krzyżak, A.; Walk, H.:* Estimation of the essential supremum of a regression function
- 2010/009 *Geisinger, L.; Laptev, A.; Weidl, T.:* Geometrical Versions of improved Berezin-Li-Yau Inequalities
- 2010/008 *Poppitz, S.; Stroppel, M.:* Polarities of Schellhammer Planes
- 2010/007 *Grundhöfer, T.; Krinn, B.; Stroppel, M.:* Non-existence of isomorphisms between certain unitals
- 2010/006 *Höllig, K.; Hörner, J.; Hoffacker, A.:* Finite Element Analysis with B-Splines: Weighted and Isogeometric Methods

- 2010/005 *Kaltenbacher, B.; Walk, H.:* On convergence of local averaging regression function estimates for the regularization of inverse problems
- 2010/004 *Kühnel, W.; Solanes, G.:* Tight surfaces with boundary
- 2010/003 *Kohler, M; Walk, H.:* On optimal exercising of American options in discrete time for stationary and ergodic data
- 2010/002 *Gulde, M.; Stoppel, M.:* Stabilizers of Subspaces under Similitudes of the Klein Quadric, and Automorphisms of Heisenberg Algebras
- 2010/001 *Leitner, F.:* Examples of almost Einstein structures on products and in cohomogeneity one
- 2009/008 *Griesemer, M.; Zenk, H.:* On the atomic photoeffect in non-relativistic QED
- 2009/007 *Griesemer, M.; Moeller, J.S.:* Bounds on the minimal energy of translation invariant n-polaron systems
- 2009/006 *Demirel, S.; Harrell II, E.M.:* On semiclassical and universal inequalities for eigenvalues of quantum graphs
- 2009/005 *Bächle, A, Kimmerle, W.:* Torsion subgroups in integral group rings of finite groups
- 2009/004 *Geisinger, L.; Weidl, T.:* Universal bounds for traces of the Dirichlet Laplace operator
- 2009/003 *Walk, H.:* Strong laws of large numbers and nonparametric estimation
- 2009/002 *Leitner, F.:* The collapsing sphere product of Poincaré-Einstein spaces
- 2009/001 *Brehm, U.; Kühnel, W.:* Lattice triangulations of E^3 and of the 3-torus
- 2008/006 *Kohler, M.; Krzyżak, A.; Walk, H.:* Upper bounds for Bermudan options on Markovian data using nonparametric regression and a reduced number of nested Monte Carlo steps
- 2008/005 *Kaltenbacher, B.; Schöpfer, F.; Schuster, T.:* Iterative methods for nonlinear ill-posed problems in Banach spaces: convergence and applications to parameter identification problems
- 2008/004 *Leitner, F.:* Conformally closed Poincaré-Einstein metrics with intersecting scale singularities
- 2008/003 *Effenberger, F.; Kühnel, W.:* Hamiltonian submanifolds of regular polytope
- 2008/002 *Hertweck, M.; Höfert, C.R.; Kimmerle, W.:* Finite groups of units and their composition factors in the integral group rings of the groups $PSL(2, q)$
- 2008/001 *Kovarik, H.; Vugalter, S.; Weidl, T.:* Two dimensional Berezin-Li-Yau inequalities with a correction term
- 2007/006 *Weidl, T.:* Improved Berezin-Li-Yau inequalities with a remainder term
- 2007/005 *Frank, R.L.; Loss, M.; Weidl, T.:* Polya's conjecture in the presence of a constant magnetic field
- 2007/004 *Ekholm, T.; Frank, R.L.; Kovarik, H.:* Eigenvalue estimates for Schrödinger operators on metric trees
- 2007/003 *Lesky, P.H.; Racke, R.:* Elastic and electro-magnetic waves in infinite waveguides
- 2007/002 *Teufel, E.:* Spherical transforms and Radon transforms in Moebius geometry
- 2007/001 *Meister, A.:* Deconvolution from Fourier-oscillating error densities under decay and smoothness restrictions