

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Homogeneous almost quaternion-Hermitian
manifolds

Andrei Moroianu, Mihaela Pilca, Uwe Semmelmann

Preprint 2012/016

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Homogeneous almost quaternion-Hermitian
manifolds

Andrei Moroianu, Mihaela Pilca, Uwe Semmelmann

Preprint 2012/016

Fachbereich Mathematik
Fakultät Mathematik und Physik
Universität Stuttgart
Pfaffenwaldring 57
D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de

WWW: <http://www.mathematik.uni-stuttgart.de/preprints>

ISSN **1613-8309**

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.
L^AT_EX-Style: Winfried Geis, Thomas Merkle

HOMOGENEOUS ALMOST QUATERNION-HERMITIAN MANIFOLDS

ABSTRACT. An almost quaternion-Hermitian structure on a Riemannian manifold (M^{4n}, g) is a reduction of the structure group of M to $\mathrm{Sp}(n)\mathrm{Sp}(1) \subset \mathrm{SO}(4n)$. In this paper we show that a compact simply connected homogeneous almost quaternion-Hermitian manifold of non-vanishing Euler characteristic is either a Wolf space, or $\mathbb{S}^2 \times \mathbb{S}^2$, or the complex quadric $\mathrm{SO}(7)/\mathrm{U}(3)$.

2010 *Mathematics Subject Classification*: Primary: 53C30, 53C35, 53C15. Secondary: 17B22

Keywords: Quaternion-Hermitian structures, homogeneous spaces, root systems, Clifford structures.

1. INTRODUCTION

The notion of (even) Clifford structures on Riemannian manifolds was introduced in [12]. Roughly speaking, a rank r (even) Clifford structure on M is a rank r Euclidean bundle whose (even) Clifford algebra bundle acts on the tangent bundle of M . For $r = 3$, an even Clifford structure on M is just an almost quaternionic structure, i.e. a rank 3 sub-bundle Q of the endomorphism bundle $\mathrm{End}(TM)$ locally spanned by three endomorphisms I, J, K satisfying the quaternionic relations

$$I^2 = J^2 = K^2 = -\mathrm{id}, \quad IJ = K.$$

If moreover $Q \subset \mathrm{End}^-(TM)$ (or, equivalently, if I, J, K are g -orthogonal), the structure (M, g, Q) is called almost quaternion-Hermitian [7, 8, 9, 17].

Homogeneous even Clifford structures on homogeneous compact manifolds of non-vanishing Euler characteristic were studied in [11], where it is established an upper bound for their rank, as well as a description of the limiting cases. In this paper we consider the other extremal case, namely even Clifford structures with the lowest possible (non-trivial) rank, which is 3 and give the complete classification of compact homogeneous almost quaternion-Hermitian manifolds G/H with non-vanishing Euler characteristic. This last assumption turns out to be crucial at several places throughout the proof (see below). Without it, the classification is completely out of reach, but there are lots of homogeneous examples constructed for instance by D. Joyce [4, 5] and O. Maciá [6].

Our classification result is the following:

Theorem 1.1. *A compact simply connected homogeneous manifold $M = G/H$ of non-vanishing Euler characteristic carries a homogeneous almost quaternion-Hermitian structure if and only if it belongs to the following list:*

This work was supported by the contract ANR-10-BLAN 0105 “Aspects Conformes de la Géométrie”. The second-named author thanks the Centre de Mathématiques de l’École Polytechnique for hospitality during the preparation of this work.

- *Wolf spaces* G/N where G is any compact simple Lie group and N is the normalizer of some subgroup $\mathrm{Sp}(1) \subset G$ determined by a highest root of G , cf. [18].
- $\mathbb{S}^2 \times \mathbb{S}^2$.
- $\mathrm{SO}(7)/\mathrm{U}(3)$.

Let us first give some comments on the above list. The Wolf spaces are quaternion-Kähler manifolds [18], so they admit not only a topological but even a *holonomy* reduction to $\mathrm{Sp}(n)\mathrm{Sp}(1)$. In dimension 4, every orientable manifold is almost quaternion-Hermitian since $\mathrm{Sp}(1)\mathrm{Sp}(1) = \mathrm{SO}(4)$. In this dimension there exist (up to homothety) only two compact simply connected homogeneous manifolds with non-vanishing Euler characteristic: $\mathbb{S}^2 \times \mathbb{S}^2$ and \mathbb{S}^4 . The latter is already a Wolf space since $\mathbb{S}^4 = \mathbb{H}\mathbb{P}^1$, this is why in dimension 4, the only extra space in the list is $\mathbb{S}^2 \times \mathbb{S}^2$. Finally, the complex quadric $\mathrm{SO}(7)/\mathrm{U}(3) \subset \mathbb{C}\mathbb{P}^7$, which incidentally is also the twistor space of \mathbb{S}^6 , carries a 1-parameter family of $\mathrm{Sp}(3)\mathrm{U}(1)$ structures with fixed volume. Motivated by our present classification, F. Martín Cabrera and A. Swann [10] are currently investigating the quaternion Hermitian type of this family.

The outline of the proof of Theorem 1.1 is as follows: The first step is to show (in Proposition 3.3) that G has to be a simple Lie group, unless $M = \mathbb{S}^2 \times \mathbb{S}^2$. The condition $\chi(M) \neq 0$ (which is equivalent to $\mathrm{rk}(H) = \mathrm{rk}(G)$) is used here in order to ensure that every subgroup of maximal rank of a product $G_1 \times G_2$ is itself a product. The next step is to rule out the case $G = \mathrm{G}_2$ which is the only simple group for which the ratio between the length of the long and short roots is $\sqrt{3}$. Once this is done, we can thus assume that either all roots of G have the same length, or the ratio between the length of the long and short roots is $\sqrt{2}$. We further show that if G/H is symmetric, then H has an $\mathrm{Sp}(1)$ -factor, so M is a Wolf space.

Now, since $\mathrm{rk}(H) = \mathrm{rk}(G)$, the weights of the (complexified) isotropy representation $\mathfrak{m}^{\mathbb{C}}$ can be identified with a subset of the root system of G . We show that the existence of a homogeneous almost quaternion-Hermitian structure on G/H implies that the set of weights $\mathcal{W}(\mathfrak{m}^{\mathbb{C}})$ can be split into two distinct subsets, one of which is obtained from the other by a translation (Proposition 3.1 below). Moreover, if G/H is not symmetric, then $[\mathfrak{m}, \mathfrak{m}] \cap \mathfrak{m} \neq 0$, so $(\mathcal{W}(\mathfrak{m}^{\mathbb{C}}) + \mathcal{W}(\mathfrak{m}^{\mathbb{C}})) \cap \mathcal{W}(\mathfrak{m}^{\mathbb{C}}) \neq \emptyset$. Putting all this information together we are then able to show, using the properties of root systems, that there is one single isotropy weight system satisfying these conditions, namely the isotropy representation of $\mathrm{SO}(7)/\mathrm{U}(3)$, whose restriction to $\mathrm{SU}(3)$ is isomorphic to $\mathbb{C}^3 \oplus (\mathbb{C}^3)^*$ and is therefore quaternionic.

2. PRELIMINARIES

Let $M = G/H$ be a homogeneous space. Throughout this paper we make the following assumptions:

- M is compact (and thus G and H are compact, too).
- The infinitesimal isotropy representation is faithful (this is always the case after taking an appropriate quotient of G)
- M has non-vanishing Euler characteristic: $\chi(M) \neq 0$, or, equivalently, $\mathrm{rk}(H) = \mathrm{rk}(G)$.

- M is simply connected. An easy argument using the exact homotopy sequence shows that by changing the representation of M as homogeneous space if necessary, one can assume that G is simply connected and H is connected (see [13] for example).

Denote by \mathfrak{h} and \mathfrak{g} the Lie algebras of H and G and by \mathfrak{m} the orthogonal complement of \mathfrak{h} in \mathfrak{g} with respect to some $\text{ad}_{\mathfrak{g}}$ -invariant scalar product on \mathfrak{g} . The restriction to \mathfrak{m} of this scalar product defines a homogeneous Riemannian metric g on M .

An almost quaternion-Hermitian structure on a Riemannian manifold (M, g) is a three-dimensional sub-bundle of the bundle of skew-symmetric endomorphisms $\text{End}^-(TM)$, which is locally spanned by three endomorphisms satisfying the quaternion relations [7, 17]. In the case where $M = G/H$ is homogeneous, such a structure is called homogeneous if this three-dimensional sub-bundle is defined by a three-dimensional H -invariant summand of the second exterior power of the isotropy representation $\Lambda^2\mathfrak{m} = \text{End}^-(\mathfrak{m})$. For our purposes, we give the following equivalent definition, which corresponds to the fact that an almost quaternion-Hermitian structure is just a rank 3 even Clifford structure (cf. [11, 12]):

Definition 2.1. A homogeneous almost quaternion-Hermitian structure on the Riemannian homogeneous space $(G/H, g)$ is an orthogonal representation $\rho : H \rightarrow \text{SO}(3)$ and an H -equivariant Lie algebra morphism $\varphi : \mathfrak{so}(3) \rightarrow \text{End}^-(\mathfrak{m})$ extending to an algebra representation of the even real Clifford algebra Cl_3^0 on \mathfrak{m} .

The H -equivariance of the morphism $\varphi : \mathfrak{so}(3) \rightarrow \text{End}^-(\mathfrak{m})$ is with respect to the following actions of H : the action on $\mathfrak{so}(3)$ is given by the composition of the adjoint representation of $\text{SO}(3)$ with ρ , and the action on $\text{End}^-(\mathfrak{m})$ is the one induced by the isotropy representation ι of H . Since φ extends to a representation of $\text{Cl}_3^0 \simeq \mathbb{H}$ on \mathfrak{m} , the above definition readily implies the following result (see also [11, Lemma 3.2] or [15]):

Lemma 2.2. The complexified isotropy representation ι_* on $\mathfrak{m}^{\mathbb{C}}$ is isomorphic to the tensor product $\mathfrak{m}^{\mathbb{C}} = \mathbb{H} \otimes_{\mathbb{C}} \mathbb{E}$, where \mathbb{H} is defined by the composition $\mu := \xi \circ \rho_*$ of ρ_* with the spin representation ξ of $\mathfrak{so}(3) = \mathfrak{spin}(3) = \mathfrak{sp}(1)$ on \mathbb{H} , and \mathbb{E} is defined by the composition $\lambda := \pi \circ \iota_*$ of the isotropy representation with the projection of \mathfrak{h} to the kernel of ρ_* .

3. THE CLASSIFICATION

In this section we classify all compact simply connected homogeneous almost quaternion-Hermitian manifolds $M = G/H$ with non-vanishing Euler characteristic.

We choose a common maximal torus of H and G and denote by $\mathfrak{t} \subset \mathfrak{h}$ its Lie algebra. Then the root system $\mathcal{R}(\mathfrak{g}) \subset \mathfrak{t}^*$ is the disjoint union of the root system $\mathcal{R}(\mathfrak{h})$ and the set \mathcal{W} of weights of the complexified isotropy representation of the homogeneous space G/H . This follows from the fact that the isotropy representation is given by the restriction to H of the adjoint representation of \mathfrak{g} .

The weights of the complex spin representation of $\mathfrak{so}(3)$ on $\Sigma_3^{\mathbb{C}} \simeq \mathbb{H}$ are $\mathcal{W}(\Sigma_3^{\mathbb{C}}) = \{\pm \frac{1}{2}e_1\}$, where e_1 is some element of norm 1 of the dual of some Cartan sub-algebra of $\mathfrak{so}(3)$. We denote by $\beta \in \mathfrak{t}^*$ the pull-back through μ of the vector $\frac{1}{2}e_1$ and by $\mathcal{A} := \{\pm\alpha_1, \dots, \pm\alpha_n\} \subset \mathfrak{t}^*$ the weights of the self-dual representation λ . By Lemma 2.2, we

obtain the following description of the weights of the isotropy representation of any homogeneous almost quaternion-Hermitian manifold $M = G/H$, which is a particular case of [11, Proposition 3.3]:

Proposition 3.1. *The set $\mathcal{W} := \mathcal{W}(\mathfrak{m})$ of weights of the isotropy representation is given by:*

$$(1) \quad \mathcal{W} = \{\varepsilon_i \alpha_i + \varepsilon \beta\}_{1 \leq i \leq n; \varepsilon_i, \varepsilon \in \{\pm 1\}}.$$

As an immediate consequence we have:

Lemma 3.2. *Let $(G/H, g, \rho, \varphi)$ be a homogeneous almost quaternion-Hermitian structure as in Definition 2.1. Then the infinitesimal representation $\rho_* : \mathfrak{h} \rightarrow \mathfrak{so}(3)$ does not vanish.*

Proof. Suppose for a contradiction that $\rho_* = 0$. Then the \mathfrak{h} -representation H defined in Lemma 2.2 is trivial, so $\beta = 0$ and $\mathfrak{m}^{\mathbb{C}} = E \oplus E$. Every weight of the (complexified) isotropy representation appears then twice in the root system of G , which is impossible (cf. [16, p. 38]). \square

Our next goal is to show that the automorphism group of a homogeneous almost quaternion-Hermitian manifold is in general a simple Lie group:

Proposition 3.3. *If G/H is a simply connected compact homogeneous almost quaternion-Hermitian manifold with non-vanishing Euler characteristic, then either G is simple or $G = \mathrm{SU}(2) \times \mathrm{SU}(2)$ and $M = \mathbb{S}^2 \times \mathbb{S}^2$.*

Proof. We already know that G is compact and simply connected. If G is not simple, then $G = G_1 \times G_2$ with $\dim(G_i) \geq 3$. Let \mathfrak{g}_i denote the Lie algebra of G_i , so that $\mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_2$. By a classical result of Borel and Siebenthal ([3, p. 210]), the Lie algebra of the subgroup H splits as $\mathfrak{h} = \mathfrak{h}_1 \oplus \mathfrak{h}_2$, where $\mathfrak{h}_i = \mathfrak{h} \cap \mathfrak{g}_i$. Correspondingly, the isotropy representation splits as $\mathfrak{m} = \mathfrak{m}_1 \oplus \mathfrak{m}_2$, where \mathfrak{m}_i is the isotropy representation of \mathfrak{h}_i in \mathfrak{g}_i .

Let $\{e_1, e_2, e_3\}$ be an orthonormal basis of $\mathfrak{so}(3)$ and let us denote by $J_i := \varphi(e_i)$, for $1 \leq i \leq 3$. The H -equivariance of φ implies that

$$(2) \quad \varphi(\rho_*(X)e_i) = [\mathrm{ad}_X, J_i], \quad \forall X \in \mathfrak{h}, 1 \leq i \leq 3.$$

We claim that the representation ρ_* does not vanish on \mathfrak{h}_1 or on \mathfrak{h}_2 . Assume for instance that $\rho_*(\mathfrak{h}_1) = 0$. We express each endomorphism J_i of $\mathfrak{m} = \mathfrak{m}_1 \oplus \mathfrak{m}_2$ as

$$J_i = \begin{pmatrix} A_i & B_i \\ C_i & D_i \end{pmatrix}.$$

For every $X \in \mathfrak{h}_1$, (2) shows that ad_X commutes with J_i . Expressing

$$\mathrm{ad}_X = \begin{pmatrix} \mathrm{ad}_X^{\mathfrak{g}_1} & 0 \\ 0 & 0 \end{pmatrix}$$

we get in particular that $\mathrm{ad}_X^{\mathfrak{g}_1} \circ B_i = 0$ for all $X \in \mathfrak{h}_1$. On the other hand, since $\mathrm{rk}(\mathfrak{h}_1) = \mathrm{rk}(\mathfrak{g}_1)$, there exists no vector in \mathfrak{m}_1 commuting with all $X \in \mathfrak{h}_1$, so $B_i = 0$ and thus $C_i = -B_i^* = 0$ for $1 \leq i \leq 3$. However, this would imply that the map $\varphi_1 : \mathfrak{so}(3) \rightarrow \mathrm{End}^-(\mathfrak{m}_1)$

given by $\varphi_1(e_i) = A_i$ for $1 \leq i \leq 3$ is a homogeneous almost quaternion-Hermitian structure on G_1/H_1 with vanishing ρ_* , which contradicts Lemma 3.2. This proves our claim.

Now, since $\rho_* : \mathfrak{h} \rightarrow \mathfrak{so}(3)$ is a Lie algebra morphism, we must have in particular

$$[\rho_*(\mathfrak{h}_1), \rho_*(\mathfrak{h}_2)] = 0.$$

By changing the orthonormal basis $\{e_1, e_2, e_3\}$ if necessary, we thus may assume that $\rho_*(\mathfrak{h}_1) = \rho_*(\mathfrak{h}_2) = \langle e_1 \rangle$. The Lie algebras \mathfrak{h}_1 and \mathfrak{h}_2 decompose as $\mathfrak{h}_i = \mathfrak{h}'_i \oplus \langle X_i \rangle$ where $\mathfrak{h}'_i := \ker(\rho_*) \cap \mathfrak{h}_i$ and $\rho_*(X_i) = e_1$ for $1 \leq i \leq 2$.

From (2), the following relations hold:

$$(3) \quad [\mathrm{ad}_{X_i}, J_2] = J_3, \quad [\mathrm{ad}_{X_i}, J_3] = -J_2, \quad 1 \leq i \leq 2.$$

Like before we can write

$$\mathrm{ad}_{X_1} = \begin{pmatrix} \mathrm{ad}_{X_1}^{\mathfrak{g}_1} & 0 \\ 0 & 0 \end{pmatrix}, \quad \mathrm{ad}_{X_2} = \begin{pmatrix} 0 & 0 \\ 0 & \mathrm{ad}_{X_2}^{\mathfrak{g}_2} \end{pmatrix},$$

so (3) implies that $A_2 = A_3 = 0$ and $D_2 = D_3 = 0$. In particular

$$-1 = J_2^2 = \begin{pmatrix} 0 & B_2 \\ C_2 & 0 \end{pmatrix}^2 = \begin{pmatrix} B_2 C_2 & 0 \\ 0 & C_2 B_2 \end{pmatrix},$$

thus showing that B_2 defines an isomorphism between \mathfrak{m}_2 and \mathfrak{m}_1 (whose inverse is $-C_2$).

On the other hand, since by (2) ad_X commutes with J_2 for all $X \in \mathfrak{h}'_1$, we obtain as before that $\mathrm{ad}_X^{\mathfrak{g}_1} \circ B_2 = 0$ for all $X \in \mathfrak{h}'_1$. Since B_2 is onto, this shows that the isotropy representation of G_1/H_1 restricted to \mathfrak{h}'_1 vanishes, so $\mathfrak{h}'_1 = 0$ and similarly $\mathfrak{h}'_2 = 0$. We therefore have $\mathfrak{h}_1 = \mathfrak{h}_2 = \mathbb{R}$, and since $\mathrm{rk}(G_i) = \mathrm{rk}(H_i) = 1$, we get $\mathfrak{g}_1 = \mathfrak{g}_2 = \mathfrak{su}(2)$. We thus have $G = \mathrm{SU}(2) \times \mathrm{SU}(2)$, and $H = \mathbb{T}^2$ is a maximal torus, so $M = \mathbb{S}^2 \times \mathbb{S}^2$. \square

We are in position to complete the proof of our main result:

Proof of Theorem 1.1. By Proposition 3.3 we may assume that G is simple. We first study the case $G = G_2$ (this is the only simple group for which the ratio between the length of long and short roots is neither 1, nor $\sqrt{2}$). The only connected subgroups of rank 2 of G_2 are $\mathrm{U}(2), \mathrm{SU}(3), \mathrm{SO}(4)$ and \mathbb{T}^2 . The spaces $G_2/\mathrm{U}(2)$ and $G_2/\mathrm{SU}(3)$ have dimension 10 and 6 respectively, therefore they can not carry almost quaternion-Hermitian structures.

The quotient $G_2/\mathrm{SO}(4)$ is a Wolf space, so it remains to study the generalized flag manifold G_2/\mathbb{T}^2 . We claim that this space has no homogeneous almost quaternion-Hermitian structure. Indeed, if this were the case, using Proposition 3.1 one could express the root system of G_2 as the disjoint union of two subsets

$$\mathcal{W}^+ := \{\varepsilon_i \alpha_i + \beta\}_{1 \leq i \leq 3; \varepsilon_i \in \{\pm 1\}}, \quad \mathcal{W}^- := \{\varepsilon_i \alpha_i - \beta\}_{1 \leq i \leq 3; \varepsilon_i \in \{\pm 1\}}$$

such that there exists some vector v ($:= 2\beta$) with $\mathcal{W}^+ = v + \mathcal{W}^-$. On the other hand, it is easy to check that there exist no such partition of $\mathcal{R}(G_2)$.

Consider now the case where $M = G/H$ is a symmetric space. If M is a Wolf space there is nothing to prove, so assume from now on that this is not the case. The Lie algebra of H can be split as $\mathfrak{h} = \ker(\rho_*) \oplus \mathfrak{h}_0$, where \mathfrak{h}_0 denotes the orthogonal complement of $\ker(\rho_*)$. Clearly \mathfrak{h}_0 is isomorphic to $\rho_*(\mathfrak{h}) \subset \mathfrak{so}(3)$ so by Lemma 3.2, $\mathfrak{h}_0 = \mathfrak{u}(1)$ or $\mathfrak{h}_0 = \mathfrak{sp}(1)$. The

latter case can not occur since our assumption that M is not a Wolf space implies that \mathfrak{h} has no $\mathfrak{sp}(1)$ -summand. We are left with the case when $\mathfrak{h} = \ker(\rho_*) \oplus \mathfrak{u}(1)$. We claim that this case can not occur either. Indeed, if such a space would carry a homogeneous almost quaternion-Hermitian structure, then the representation of $\ker(\rho_*)$ on \mathfrak{m} would be quaternionic. Two anti-commuting complex structures I, J of \mathfrak{m} induce non-vanishing elements a_I, a_J in the center of $\ker(\rho_*)$ (see the proof of [14, Lemma 2.4]). On the other hand, the adjoint actions of a_I and a_J on \mathfrak{m} are proportional to I and J respectively ([14, Eq. (4)]) and thus anti-commute, contradicting the fact that a_I and a_J commute (being central elements).

We can assume from now on, that $M = G/H$ is non-symmetric, G is simple and $G \neq G_2$. Up to a rescaling of the ad_G -invariant metric on \mathfrak{g} , we may thus assume that all roots of \mathfrak{g} have square length equal to 1 or 2.

From (1), it follows that

$$\mathcal{R}(\mathfrak{g}) = \mathcal{W}(\mathfrak{m}) \cup \mathcal{R}(\mathfrak{h}) = \{\varepsilon_i \alpha_i + \varepsilon \beta\}_{1 \leq i \leq n; \varepsilon_i, \varepsilon \in \{\pm 1\}} \cup \mathcal{R}(\mathfrak{h}).$$

Up to a change of signs of the α_i 's, we may assume:

$$(4) \quad \langle \beta, \alpha_i \rangle \geq 0, \quad \text{for all } 1 \leq i \leq n.$$

Then either the roots $\beta + \alpha_i$ and $\beta - \alpha_i$ of G have the same length, or $|\beta + \alpha_i|^2 = 2$ and $|\beta - \alpha_i|^2 = 1$. This shows that for each $1 \leq i \leq n$,

$$(5) \quad \langle \beta, \alpha_i \rangle \in \left\{ 0, \frac{1}{4} \right\}.$$

From the general property of root systems (10) below, it follows that:

$$(6) \quad |\beta|^2 \in \left\{ \frac{1}{4}, \frac{3}{4}, \frac{5}{4} \right\}.$$

Since the homogeneous space G/H is not symmetric, we have $[\mathfrak{m}, \mathfrak{m}] \not\subseteq \mathfrak{h}$, so there exist subscripts $i, j, k \in \{1, \dots, n\}$ such that $(\pm\beta \pm \alpha_i) + (\pm\beta \pm \alpha_j) = \pm\beta \pm \alpha_k$. Taking (5) into account, we need to check the following possible cases (up to a permutation of the subscripts):

- a) $\beta = \pm 2\alpha_1 \pm \alpha_2$.
- b) $\beta = \frac{\alpha_1}{3}$.
- c) $\beta = \frac{\alpha_1 \pm \alpha_2 \pm \alpha_3}{3}$.
- d) $\beta = \alpha_1 \pm \alpha_2 \pm \alpha_3$.

We will show that cases a), b) and c) can not occur and that in case d) there is only one solution.

- a) If $\beta = 2\alpha_1 + \alpha_2$, then $\beta + \alpha_2 = 2(\beta - \alpha_1)$ and this would imply the existence of two proportional roots, $\beta + \alpha_2$ and $\beta - \alpha_1$, in $\mathcal{W} \subseteq \mathcal{R}(\mathfrak{g})$, contradicting the property R2 of root systems (*cf.* Definition A.1). For all the other possible choices of signs in a) we obtain a similar contradiction.
- b) If $\beta = \frac{\alpha_1}{3}$, then there exist two proportional roots: $\beta + \alpha_1 = -2(\beta - \alpha_1)$ in $\mathcal{R}(\mathfrak{g})$, which again contradicts R2.

c) If $\beta = \frac{\alpha_1 \pm \alpha_2 \pm \alpha_3}{3}$, then $|\beta|^2 = \frac{1}{3}(\langle \beta, \alpha_1 \rangle \pm \langle \beta, \alpha_2 \rangle \pm \langle \beta, \alpha_3 \rangle)$. From (5) and (6), it follows that the only possibility is:

$$\beta = \frac{\alpha_1 + \alpha_2 + \alpha_3}{3}, |\beta|^2 = \frac{1}{4} \text{ and } \langle \beta, \alpha_i \rangle = \frac{1}{4}, \text{ for } 1 \leq i \leq 3.$$

Together with (10), this implies that for each $1 \leq i \leq 3$ we have: $|\beta + \alpha_i|^2 = 2$, $|\beta - \alpha_i|^2 = 1$ and $|\alpha_i|^2 = \frac{5}{4}$. Thus, for all $1 \leq i, j \leq 3$, $i \neq j$, we have:

$$\langle \beta + \alpha_i, \beta - \alpha_j \rangle = \frac{1}{4} - \langle \alpha_i, \alpha_j \rangle,$$

which by (10) must be equal to 0 or ± 1 , showing that $\langle \alpha_i, \alpha_j \rangle \in \{-\frac{3}{4}, \frac{1}{4}, \frac{5}{4}\}$. On the other hand, a direct computation shows that

$$\langle \alpha_1, \alpha_2 \rangle + \langle \alpha_1, \alpha_3 \rangle + \langle \alpha_2, \alpha_3 \rangle = \frac{1}{2} \left(9|\beta|^2 - \frac{15}{4} \right) = -\frac{3}{4},$$

which is not possible for any of the above values of the scalar products, yielding a contradiction.

d) From (5) and (6), it follows that there are three possible sub-cases:

Case 1. $\beta = \alpha_1 \pm \alpha_2 \pm \alpha_3$, $|\beta|^2 = \frac{1}{4}$, $\langle \beta, \alpha_1 \rangle = \frac{1}{4}$, $\langle \beta, \alpha_2 \rangle = \langle \beta, \alpha_3 \rangle = 0$.

Case 2. $\beta = \alpha_1 + \alpha_2 - \alpha_3$, $|\beta|^2 = \frac{1}{4}$, $\langle \beta, \alpha_i \rangle = \frac{1}{4}$, $1 \leq i \leq 3$.

Case 3. $\beta = \alpha_1 + \alpha_2 + \alpha_3$, $|\beta|^2 = \frac{3}{4}$, $\langle \beta, \alpha_i \rangle = \frac{1}{4}$, $1 \leq i \leq 3$.

Case 1. From (10) it follows $|\alpha_1|^2 = \frac{5}{4}$ and $|\alpha_2|^2 = |\alpha_3|^2 = \frac{3}{4}$. Since $\langle \beta + \alpha_1, \beta - \alpha_1 \rangle = -1$ and $|\beta + \alpha_1|^2 = 2|\beta - \alpha_1|^2$, the reflexion property (11) shows that $2\beta = (\beta + \alpha_1) + (\beta - \alpha_1)$ and $3\beta - \alpha_1 = (\beta + \alpha_1) + 2(\beta - \alpha_1)$ belong to $\mathcal{R}(\mathfrak{g})$. We show that these roots actually belong to $\mathcal{R}(\mathfrak{h})$, i.e. that $2\beta, 3\beta - \alpha_1 \notin \mathcal{W}$. We argue by contradiction.

Let us first assume that $2\beta \in \mathcal{W}$. Then there exists k , $1 \leq k \leq n$, such that $2\beta = \pm\beta \pm \alpha_k$. If $\beta = \pm\alpha_k$ we obtain that $0 = \beta \mp \alpha_k$ belongs to $\mathcal{R}(\mathfrak{g})$, which contradicts the property R1 of root systems. If $\beta = \pm\frac{\alpha_k}{3}$, then the roots $\beta + \alpha_k$ and $\beta - \alpha_k$ are proportional, which contradicts R2.

Now we assume that $3\beta - \alpha_1 \in \mathcal{W}$ and conclude similarly. In this case there exists k , $1 \leq k \leq n$ such that either $2\beta = \alpha_1 \pm \alpha_k$ or $4\beta = \alpha_1 \pm \alpha_k$. In the first case we obtain $\beta - \alpha_1 = -\beta \pm \alpha_k$, which contradicts the fact that roots of G are simple. In the second case (5) yields $|\beta|^2 = \frac{1}{4}\langle \beta, \alpha_1 \rangle \pm \frac{1}{4}\langle \beta, \alpha_k \rangle \leq \frac{1}{8}$, which contradicts (6).

This shows that $2\beta, 3\beta - \alpha_1 \in \mathcal{R}(\mathfrak{h})$. Moreover $\langle 2\beta, 3\beta - \alpha_1 \rangle = 1$ and thus, by (11), their difference is a root of \mathfrak{h} too: $\beta - \alpha_1 = (3\beta - \alpha_1) - (2\beta) \in \mathcal{R}(\mathfrak{h})$, which is in contradiction with $\beta - \alpha_1 \in \mathcal{W}$. Consequently, case 1. can not occur.

Case 2. From (10) it follows that $|\alpha_i|^2 = \frac{5}{4}$, for all $1 \leq i \leq 3$. For all $1 \leq i, j \leq 3$, $i \neq j$, we then compute: $\langle \beta + \alpha_i, \beta + \alpha_j \rangle = \frac{3}{4} + \langle \alpha_i, \alpha_j \rangle$, which by (10) must be equal to 0 or ± 1 , implying that $\langle \alpha_i, \alpha_j \rangle \in \{-\frac{7}{4}, -\frac{3}{4}, \frac{1}{4}\}$. On the other hand, we obtain

$$\langle \alpha_1, \alpha_2 \rangle + \langle \alpha_1, \alpha_3 \rangle + \langle \alpha_2, \alpha_3 \rangle = \frac{1}{2} \left(|\beta|^2 - \frac{15}{4} \right) = -\frac{7}{4},$$

which is not possible for any of the above values of the scalar products, yielding again a contradiction.

Case 3. From (10) it follows that $|\alpha_i|^2 = \frac{3}{4}$, for all $1 \leq i \leq 3$. Computing the norm of $\beta - \alpha_k = \alpha_i + \alpha_j$, where $\{i, j, k\}$ is any permutation of $\{1, 2, 3\}$, yields that $\langle \alpha_i, \alpha_j \rangle = -\frac{1}{4}$, for all $1 \leq i, j \leq 3, i \neq j$. We then get

$$\langle \beta + \alpha_i, \beta + \alpha_j \rangle = 1, \quad \text{for all } 1 \leq i, j \leq 3, i \neq j,$$

which by the reflexion property (11) implies that

$$(7) \quad \{\alpha_i - \alpha_j\}_{1 \leq i, j \leq 3} \subseteq \mathcal{R}(\mathfrak{g}).$$

We claim that $n = 3$ (recall that n denotes the number of vectors α_i , or equivalently the quaternionic dimension of M). Assume for a contradiction that $n \geq 4$. By (5), $\langle \beta, \alpha_l \rangle = \frac{1}{4}$ or $\langle \beta, \alpha_l \rangle = 0$, for any $4 \leq l \leq n$.

If $\langle \beta, \alpha_l \rangle = \frac{1}{4}$ for some $l \geq 4$, it follows that $|\alpha_l|^2 = \frac{3}{4}$ and $|\beta + \alpha_l|^2 = 2$, implying by (10) that the scalar product $\langle \beta - \alpha_i, \beta + \alpha_l \rangle$ belongs to $\{\pm 1, 0\}$, for $1 \leq i \leq 3$. This further yields that $\langle \alpha_i, \alpha_l \rangle \in \{\frac{7}{4}, \frac{3}{4}, -\frac{1}{4}\}$. On the other hand, the Cauchy-Schwarz inequality applied to α_i and α_l and the fact that \mathcal{W} has only simple roots (being a root sub-system) imply that the only possible value is $\langle \alpha_i, \alpha_l \rangle = -\frac{1}{4}$, for $1 \leq i \leq 3$ and $4 \leq l \leq n$. Thus, $|\beta + \alpha_l|^2 = 0$, which contradicts the property R1 of root systems (*cf.* Definition A.1).

We therefore have $\langle \beta, \alpha_l \rangle = 0$, for all $4 \leq l \leq n$. If $|\beta \pm \alpha_l|^2 = 2$ for some $l \geq 4$ then $|\alpha_l|^2 = \frac{5}{4}$, so $\langle \beta - \alpha_l, \beta + \alpha_l \rangle = -\frac{1}{2}$, contradicting (10). Thus $|\beta \pm \alpha_l|^2 = 1$ for all $4 \leq l \leq n$. If $n \geq 5$, (10) implies

$$\langle \beta - \alpha_l, \beta + \alpha_s \rangle, \langle \beta - \alpha_l, \beta - \alpha_s \rangle \in \left\{0, \pm \frac{1}{2}\right\}, \quad \text{for } 4 \leq l, s \leq n, l \neq s.$$

This contradicts the equality $\langle \beta - \alpha_l, \beta + \alpha_s \rangle + \langle \beta - \alpha_l, \beta - \alpha_s \rangle = \frac{3}{2}$, showing that $n \leq 4$.

It remains to show that the existence of $\alpha_4 \in \mathcal{A}$, which by the above necessarily satisfies $\langle \beta, \alpha_4 \rangle = 0$ and $|\alpha_4|^2 = \frac{1}{4}$, leads to a contradiction. By (10), it follows that

$$1 + \langle \alpha_i, \alpha_4 \rangle = \langle \beta + \alpha_i, \beta + \alpha_4 \rangle \in \{\pm 1, 0\}, \quad \forall 1 \leq i \leq 3.$$

This constraint together with the Cauchy-Schwarz inequality, $|\langle \alpha_i, \alpha_4 \rangle| \leq \frac{\sqrt{3}}{4}$, implies that $\langle \alpha_i, \alpha_4 \rangle = 0$, for $1 \leq i \leq 3$.

Applying the reflexion property (11) to $\beta + \alpha_4$ and $\beta + \alpha_i$, for $1 \leq i \leq 3$, which satisfy $\langle \beta + \alpha_i, \beta + \alpha_4 \rangle = 1$ and $|\beta + \alpha_i|^2 = 2|\beta + \alpha_4|^2$, it follows that $\alpha_i - \alpha_4, \beta + 2\alpha_4 - \alpha_i \in \mathcal{R}(\mathfrak{g})$. We now show that all these roots actually belong to $\mathcal{R}(\mathfrak{h})$. Let us assume that $\alpha_i - \alpha_4 \in \mathcal{W}$ for some $i \leq 3$, i.e. there exists $s, 1 \leq s \leq 4$, such that $\alpha_i - \alpha_4 \in \{\pm\beta \pm \alpha_s\}$. Since α_4 is orthogonal to β and to α_i , for $1 \leq i \leq 3$, it follows that $\alpha_i - \alpha_4$ must be equal to $\pm\beta - \alpha_4$, leading to the contradiction that $0 = \beta \mp \alpha_i \in \mathcal{W}$. Therefore $\alpha_i - \alpha_4 \in \mathcal{R}(\mathfrak{h})$. A similar argument shows that $\beta + 2\alpha_4 - \alpha_i \in \mathcal{R}(\mathfrak{h})$.

Now, since the scalar product of these two roots of \mathfrak{h} is $\langle \alpha_i - \alpha_4, \beta + 2\alpha_4 - \alpha_i \rangle = -1$, it follows again by (11) that their sum $\beta + \alpha_4$ also belongs to $\mathcal{R}(\mathfrak{h})$, contradicting the fact that $\beta + \alpha_4 \in \mathcal{W}(\mathfrak{m})$. This finishes the proof of the claim that $n = 3$.

Since the determinant of the Gram matrix $(\langle \alpha_i, \alpha_j \rangle)_{1 \leq i, j \leq 3}$ is equal to $\frac{5}{16}$, the vectors $\{\alpha_i\}_{1 \leq i \leq 3}$ are linearly independent. Thus the roots of \mathfrak{g} given by (7) can not belong to \mathcal{W} , and therefore $\{\alpha_i - \alpha_j\}_{1 \leq i, j \leq 3}$ belong to $\mathcal{R}(\mathfrak{h})$.

Concluding, we have proven that $n = 3$ and that the following inclusions hold (after introducing the notation $\gamma_i := \alpha_j + \alpha_k$ for all permutations $\{i, j, k\}$ of $\{1, 2, 3\}$):

$$(8) \quad \{\gamma_i - \gamma_j\}_{1 \leq i \neq j \leq 3} \subseteq \mathcal{R}(\mathfrak{h}), \quad \{\gamma_i - \gamma_j\}_{1 \leq i \neq j \leq 3} \cup \{\pm \gamma_i\}_{1 \leq i \leq 3} \subseteq \mathcal{R}(\mathfrak{g}),$$

where $\langle \gamma_i, \gamma_j \rangle = \delta_{ij}$, for all $1 \leq i, j \leq 3$.

Since these sets are closed root systems and we are interested in the representation of M as a homogeneous space G/H with the smallest possible group G , we may assume that we have equality in (8). Hence $\mathcal{R}(\mathfrak{h}) = \{\gamma_i - \gamma_j\}_{1 \leq i \neq j \leq 3}$, with $\{\gamma_i\}_{1 \leq i \leq 3}$ an orthonormal basis, (which is exactly the root system of the Lie algebra $\mathfrak{su}(3)$), and $\mathcal{R}(\mathfrak{g}) = \{\gamma_i - \gamma_j\}_{1 \leq i \neq j \leq 3} \cup \{\pm \gamma_i\}_{1 \leq i \leq 3}$, which is the root system of $\mathfrak{so}(7)$. We conclude that the only possible solution is the simply connected homogeneous space $\text{SO}(7)/\text{U}(3)$.

It remains to check that this space indeed carries a homogeneous almost quaternionic-Hermitian structure. Using the sequence of inclusions

$$\mathfrak{u}(3) \subset \mathfrak{so}(6) \subset \mathfrak{so}(7),$$

we see that the isotropy representation \mathfrak{m} of $\text{SO}(7)/\text{U}(3)$ is the direct sum of the restriction to $\text{U}(3)$ of the isotropy representation of the sphere $\text{SO}(7)/\text{SO}(6)$, (which is just the standard representation of $\text{U}(3)$ on \mathbb{C}^3), and of the isotropy representation of $\text{SO}(6)/\text{U}(3)$, which is $\Lambda^2(\mathbb{C}^3)$ (cf. [2, p. 312]):

$$\mathfrak{m} = \mathbb{C}^3 \oplus \Lambda^2(\mathbb{C}^3).$$

Let I denote the complex structure of \mathfrak{m} . After identifying $\text{U}(1)$ with the center of $\text{U}(3)$ via the diagonal embedding, an element $z \in \text{U}(1)$ acts on \mathfrak{m} by complex multiplication with z^3 , i.e. $\iota(z) = z^3$. Since $\Lambda^2(\mathbb{C}^3) = (\mathbb{C}^3)^*$ as complex $\text{SU}(3)$ -representations, it follows that the restriction to $\text{SU}(3)$ of the isotropy representation \mathfrak{m} is $\mathbb{C}^3 \oplus (\mathbb{C}^3)^*$, and thus carries a quaternionic structure, i.e. a complex anti-linear automorphism J . We claim that a homogeneous almost quaternionic-Hermitian structure on $\text{SO}(7)/\text{U}(3)$ in the sense of Definition 2.1 is given by $\rho : \text{U}(3) \rightarrow \text{SO}(3)$ and $\varphi : \mathfrak{so}(3) \simeq \text{Im}(\mathbb{H}) \rightarrow \text{End}^-(\mathfrak{m})$ defined by

$$\rho(A) = \det(A), \quad \varphi(i) = I, \quad \varphi(j) = J, \quad \varphi(k) = IJ,$$

where $\det(A) \in \text{U}(1)$ is viewed as an element in $\text{SO}(3)$ via the composition

$$\text{U}(1) = S(\mathbb{C}) \rightarrow S(\mathbb{H}) = \text{Spin}(3) \rightarrow \text{SO}(3).$$

Indeed, the only thing to check is the equivariance of φ , i.e.

$$(9) \quad \varphi(\rho(A)M\rho(A)^{-1}) = \iota(A)\varphi(M)\iota(A)^{-1}, \quad \forall M \in \mathfrak{so}(3), \quad \forall A \in \text{U}(3).$$

Write $A = zB$ with $B \in \text{SU}(3)$. Then $\rho(A) = z^3$, $\iota(A) = z^3\iota(B)$ and $\iota(B)$ commutes with I, J, K , thus with $\varphi(M)$. The relation (9) is trivially satisfied for $M = i$, whereas for $M = j$ or $M = k$ one has $Mz = \bar{z}M = z^{-1}M$ and similarly $\varphi(M)\iota(z)\iota(B) = \iota(z^{-1})\iota(B)\varphi(M)$, so

$$\varphi(\rho(A)M\rho(A)^{-1}) = \varphi(z^3 M z^{-3}) = \varphi(z^6 M) = z^6 \varphi(M) = \iota(z^2) \varphi(M) = \iota(A) \varphi(M) \iota(A)^{-1}.$$

This finishes the proof of the theorem. \square

APPENDIX A. ROOT SYSTEMS

For the basic theory of root systems we refer to [1] and [16].

Definition A.1. A set \mathcal{R} of vectors in a Euclidean space $(V, \langle \cdot, \cdot \rangle)$ is called a *root system* if it satisfies the following conditions:

R1: \mathcal{R} is finite, $\text{span}(\mathcal{R}) = V$, $0 \notin \mathcal{R}$.

R2: If $\alpha \in \mathcal{R}$, then the only multiples of α in \mathcal{R} are $\pm\alpha$.

R3: $\frac{2\langle \alpha, \beta \rangle}{\langle \alpha, \alpha \rangle} \in \mathbb{Z}$, for all $\alpha, \beta \in \mathcal{R}$.

R4: $s_\alpha : \mathcal{R} \rightarrow \mathcal{R}$, for all $\alpha \in \mathcal{R}$ (s_α is the reflection $s_\alpha : V \rightarrow V$, $s_\alpha(v) := v - \frac{2\langle \alpha, v \rangle}{\langle \alpha, \alpha \rangle} \alpha$).

Let G be a compact semi-simple Lie group with Lie algebra \mathfrak{g} endowed with an $\text{ad}_{\mathfrak{g}}$ -invariant scalar product. Fix a Cartan sub-algebra $\mathfrak{t} \subset \mathfrak{g}$ and let $\mathcal{R}(\mathfrak{g}) \subset \mathfrak{t}^*$ denote its root system. It is well-known that $\mathcal{R}(\mathfrak{g})$ satisfies the conditions in Definition A.1. Conversely, every set of vectors satisfying the conditions in Definition A.1 is the root system of a unique semi-simple Lie algebra of compact type.

Remark A.2 (Properties of root systems). Let \mathcal{R} be a root system. If $\alpha, \beta \in \mathcal{R}$ such that $\beta \neq \pm\alpha$ and $\|\beta\|^2 \geq \|\alpha\|^2$, then either $\langle \alpha, \beta \rangle = 0$ or

$$(10) \quad \left(\frac{\|\beta\|^2}{\|\alpha\|^2}, \frac{2\langle \alpha, \beta \rangle}{\langle \alpha, \alpha \rangle} \right) \in \{(1, \pm 1), (2, \pm 2), (3, \pm 3)\}.$$

In other words, either the scalar product of two roots vanishes, or its absolute value equals half the square length of the longest root. Moreover,

$$(11) \quad \beta - \text{sgn} \left(\frac{2\langle \alpha, \beta \rangle}{\langle \alpha, \alpha \rangle} \right) k\alpha \in \mathcal{R}, \quad \text{for all } k \in \mathbb{Z}, 1 \leq k \leq \left\lfloor \frac{2\langle \alpha, \beta \rangle}{\langle \alpha, \alpha \rangle} \right\rfloor.$$

Definition A.3 ([11]). A set \mathcal{P} of vectors in a Euclidean space $(V, \langle \cdot, \cdot \rangle)$ is called a *root sub-system* if it satisfies the conditions R1 - R3 from Definition A.1 and if the set $\overline{\mathcal{P}}$ obtained from \mathcal{P} by taking all possible reflections is a root system.

REFERENCES

- [1] J. Adams, *Lectures on Lie groups*, The University of Chicago Press, Chicago (1969).
- [2] A. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **10** Springer-Verlag, Berlin, 1987.
- [3] A. Borel, J. de Siebenthal, *Les sous-groupes fermés de rang maximum des groupes de Lie clos*, Comment. Math. Helv. **23** (1949), 200–221.
- [4] D. Joyce, *The hypercomplex quotient and the quaternionic quotient*, Math. Ann. **290** (1991), 323–340.
- [5] D. Joyce, *Compact hypercomplex and quaternionic manifolds*, J. Differ. Geom. **35** (1992), 743–761.
- [6] O. Maciá, *A nearly quaternionic structure on SU(3)*, J. Geom. Phys. **60** (2010), no. 5, 791–798.
- [7] F. Martín Cabrera, *Almost Quaternion-Hermitian Manifolds*, Ann. Global Anal. Geom. **25** (2004), 277–301.
- [8] F. Martín Cabrera, A. Swann, *Almost Hermitian structures and quaternionic geometries*, Differential Geom. Appl. **21** (2004), no. 2, 199–214.

- [9] F. Martín Cabrera, A. Swann, *The intrinsic torsion of almost quaternion-Hermitian manifolds*, Ann. Inst. Fourier **58** (2008), no. 5, 1455–1497.
- [10] F. Martín Cabrera, A. Swann, *Quaternion Geometries on the Twistor Space of the Six-Sphere*, in preparation.
- [11] A. Moroianu, M. Pilca, *Higher Rank Homogeneous Clifford structures*, to appear in J. London Math. Soc.
- [12] A. Moroianu, U. Semmelmann, *Clifford structures on Riemannian manifolds*, Adv. Math. **228** (2011), 940–967.
- [13] A. Moroianu, U. Semmelmann, *Weakly complex homogeneous spaces*, to appear in J. reine angew. Math. doi:10.1515/crelle-2012-0077.
- [14] A. Moroianu, U. Semmelmann, *Invariant four-forms and symmetric pairs*, to appear in Ann. Global Anal. Geom. doi:10.1007/s10455-012-9336-y.
- [15] S. M. Salamon, *Quaternionic Kähler manifolds*, Invent. Math. **67** (1982), 143–171.
- [16] H. Samelson, *Notes on Lie Algebras*, Springer-Verlag (1990).
- [17] A. F. Swann, *Some Remarks on Quaternion-Hermitian Manifolds*, Arch. Math. **33** (1997), 349–354.
- [18] J. A. Wolf, *Complex homogeneous contact manifolds and quaternionic symmetric spaces*, J. Math. Mech. **14** (1965), 1033–1047.

Andrei Moroianu
Université de Versailles-St Quentin
Laboratoire de Mathématiques
UMR 8100 du CNRS
45 avenue des États-Unis
78035 Versailles *and* CMLS
École Polytechnique
UMR 7640 du CNRS
91128 Palaiseau, France
E-Mail: am@math.polytechnique.fr

Mihaela Pilca
Fakultät für Mathematik
Universität Regensburg
Universitätsstr. 31 D-93040 Regensburg, Germany *and* Institute of Mathematics “Simion Stoilow” of the
Romanian Academy, 21, Calea Grivitei Str. 010702-Bucharest, Romania
E-Mail: Mihaela.Pilca@mathematik.uni-regensburg.de

Uwe Semmelmann
Institut für Geometrie und Topologie
Fachbereich Mathematik
Universität Stuttgart
Pfaffenwaldring 57
70569 Stuttgart, Germany
E-Mail: uwe.semmelmann@mathematik.uni-stuttgart.de

Erschienenene Preprints ab Nummer 2007/2007-001

Komplette Liste: <http://www.mathematik.uni-stuttgart.de/preprints>

- 2012-016 *Moroianu, A.; Pilca, M.; Semmelmann, U.:* Homogeneous almost quaternion-Hermitian manifolds
- 2012-015 *Steinke, G.F.; Stroppel, M.J.:* Simple groups acting two-transitively on the set of generators of a finite elation Laguerre plane
- 2012-014 *Steinke, G.F.; Stroppel, M.J.:* Finite elation Laguerre planes admitting a two-transitive group on their set of generators
- 2012-013 *Diaz Ramos, J.C.; Dominguez Vázquez, M.; Kollross, A.:* Polar actions on complex hyperbolic spaces
- 2012-012 *Moroianu, A.; Semmelmann, U.:* Weakly complex homogeneous spaces
- 2012-011 *Moroianu, A.; Semmelmann, U.:* Invariant four-forms and symmetric pairs
- 2012-010 *Hamilton, M.J.D.:* The closure of the symplectic cone of elliptic surfaces
- 2012-009 *Hamilton, M.J.D.:* Iterated fibre sums of algebraic Lefschetz fibrations
- 2012-008 *Hamilton, M.J.D.:* The minimal genus problem for elliptic surfaces
- 2012-007 *Ferrario, P.:* Partitioning estimation of local variance based on nearest neighbors under censoring
- 2012-006 *Stroppel, M.:* Buttons, Holes and Loops of String: Lacing the Doily
- 2012-005 *Hantsch, F.:* Existence of Minimizers in Restricted Hartree-Fock Theory
- 2012-004 *Grundhöfer, T.; Stroppel, M.; Van Maldeghem, H.:* Unitals admitting all translations
- 2012-003 *Hamilton, M.J.D.:* Representing homology classes by symplectic surfaces
- 2012-002 *Hamilton, M.J.D.:* On certain exotic 4-manifolds of Akhmedov and Park
- 2012-001 *Jentsch, T.:* Parallel submanifolds of the real 2-Grassmannian
- 2011-028 *Spreer, J.:* Combinatorial 3-manifolds with cyclic automorphism group
- 2011-027 *Griesemer, M.; Hantsch, F.; Wellig, D.:* On the Magnetic Pekar Functional and the Existence of Bipolarons
- 2011-026 *Müller, S.:* Bootstrapping for Bandwidth Selection in Functional Data Regression
- 2011-025 *Felber, T.; Jones, D.; Kohler, M.; Walk, H.:* Weakly universally consistent static forecasting of stationary and ergodic time series via local averaging and least squares estimates
- 2011-024 *Jones, D.; Kohler, M.; Walk, H.:* Weakly universally consistent forecasting of stationary and ergodic time series
- 2011-023 *Györfi, L.; Walk, H.:* Strongly consistent nonparametric tests of conditional independence
- 2011-022 *Ferrario, P.G.; Walk, H.:* Nonparametric partitioning estimation of residual and local variance based on first and second nearest neighbors
- 2011-021 *Eberts, M.; Steinwart, I.:* Optimal regression rates for SVMs using Gaussian kernels
- 2011-020 *Frank, R.L.; Geisinger, L.:* Refined Semiclassical Asymptotics for Fractional Powers of the Laplace Operator
- 2011-019 *Frank, R.L.; Geisinger, L.:* Two-term spectral asymptotics for the Dirichlet Laplacian on a bounded domain
- 2011-018 *Hänel, A.; Schulz, C.; Wirth, J.:* Embedded eigenvalues for the elastic strip with cracks
- 2011-017 *Wirth, J.:* Thermo-elasticity for anisotropic media in higher dimensions
- 2011-016 *Höllig, K.; Hörner, J.:* Programming Multigrid Methods with B-Splines
- 2011-015 *Ferrario, P.:* Nonparametric Local Averaging Estimation of the Local Variance Function
- 2011-014 *Müller, S.; Dippon, J.:* k-NN Kernel Estimate for Nonparametric Functional Regression in Time Series Analysis
- 2011-013 *Knarr, N.; Stroppel, M.:* Unitals over composition algebras
- 2011-012 *Knarr, N.; Stroppel, M.:* Baer involutions and polarities in Moufang planes of characteristic two
- 2011-011 *Knarr, N.; Stroppel, M.:* Polarities and planar collineations of Moufang planes

- 2011-010 *Jentsch, T.; Moroianu, A.; Semmelmann, U.:* Extrinsic hyperspheres in manifolds with special holonomy
- 2011-009 *Wirth, J.:* Asymptotic Behaviour of Solutions to Hyperbolic Partial Differential Equations
- 2011-008 *Stroppel, M.:* Orthogonal polar spaces and unitals
- 2011-007 *Nagl, M.:* Charakterisierung der Symmetrischen Gruppen durch ihre komplexe Gruppenalgebra
- 2011-006 *Solanes, G.; Teufel, E.:* Horo-tightness and total (absolute) curvatures in hyperbolic spaces
- 2011-005 *Ginoux, N.; Semmelmann, U.:* Imaginary Kählerian Killing spinors I
- 2011-004 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic D -Scales: Part II — Gain-Scheduled Control
- 2011-003 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic D -Scales: Part I — Robust Control
- 2011-002 *Alexandrov, B.; Semmelmann, U.:* Deformations of nearly parallel G_2 -structures
- 2011-001 *Geisinger, L.; Weidl, T.:* Sharp spectral estimates in domains of infinite volume
- 2010-018 *Kimmerle, W.; Konovalov, A.:* On integral-like units of modular group rings
- 2010-017 *Gauduchon, P.; Moroianu, A.; Semmelmann, U.:* Almost complex structures on quaternion-Kähler manifolds and inner symmetric spaces
- 2010-016 *Moroianu, A.; Semmelmann, U.:* Clifford structures on Riemannian manifolds
- 2010-015 *Grafarend, E.W.; Kühnel, W.:* A minimal atlas for the rotation group $SO(3)$
- 2010-014 *Weidl, T.:* Semiclassical Spectral Bounds and Beyond
- 2010-013 *Stroppel, M.:* Early explicit examples of non-desarguesian plane geometries
- 2010-012 *Effenberger, F.:* Stacked polytopes and tight triangulations of manifolds
- 2010-011 *Györfi, L.; Walk, H.:* Empirical portfolio selection strategies with proportional transaction costs
- 2010-010 *Kohler, M.; Krzyżak, A.; Walk, H.:* Estimation of the essential supremum of a regression function
- 2010-009 *Geisinger, L.; Laptev, A.; Weidl, T.:* Geometrical Versions of improved Berezin-Li-Yau Inequalities
- 2010-008 *Poppitz, S.; Stroppel, M.:* Polarities of Schellhammer Planes
- 2010-007 *Grundhöfer, T.; Krinn, B.; Stroppel, M.:* Non-existence of isomorphisms between certain unitals
- 2010-006 *Höllig, K.; Hörner, J.; Hoffacker, A.:* Finite Element Analysis with B-Splines: Weighted and Isogeometric Methods
- 2010-005 *Kaltenbacher, B.; Walk, H.:* On convergence of local averaging regression function estimates for the regularization of inverse problems
- 2010-004 *Kühnel, W.; Solanes, G.:* Tight surfaces with boundary
- 2010-003 *Kohler, M.; Walk, H.:* On optimal exercising of American options in discrete time for stationary and ergodic data
- 2010-002 *Gulde, M.; Stroppel, M.:* Stabilizers of Subspaces under Similitudes of the Klein Quadric, and Automorphisms of Heisenberg Algebras
- 2010-001 *Leitner, F.:* Examples of almost Einstein structures on products and in cohomogeneity one
- 2009-008 *Griesemer, M.; Zenk, H.:* On the atomic photoeffect in non-relativistic QED
- 2009-007 *Griesemer, M.; Moeller, J.S.:* Bounds on the minimal energy of translation invariant n -polaron systems
- 2009-006 *Demirel, S.; Harrell II, E.M.:* On semiclassical and universal inequalities for eigenvalues of quantum graphs
- 2009-005 *Bächle, A.; Kimmerle, W.:* Torsion subgroups in integral group rings of finite groups
- 2009-004 *Geisinger, L.; Weidl, T.:* Universal bounds for traces of the Dirichlet Laplace operator
- 2009-003 *Walk, H.:* Strong laws of large numbers and nonparametric estimation
- 2009-002 *Leitner, F.:* The collapsing sphere product of Poincaré-Einstein spaces

- 2009-001 *Brehm, U.; Kühnel, W.:* Lattice triangulations of E^3 and of the 3-torus
- 2008-006 *Kohler, M.; Krzyżak, A.; Walk, H.:* Upper bounds for Bermudan options on Markovian data using nonparametric regression and a reduced number of nested Monte Carlo steps
- 2008-005 *Kaltenbacher, B.; Schöpfer, F.; Schuster, T.:* Iterative methods for nonlinear ill-posed problems in Banach spaces: convergence and applications to parameter identification problems
- 2008-004 *Leitner, F.:* Conformally closed Poincaré-Einstein metrics with intersecting scale singularities
- 2008-003 *Effenberger, F.; Kühnel, W.:* Hamiltonian submanifolds of regular polytope
- 2008-002 *Hertweck, M.; Höfert, C.R.; Kimmerle, W.:* Finite groups of units and their composition factors in the integral group rings of the groups $PSL(2, q)$
- 2008-001 *Kovarik, H.; Vugalter, S.; Weidl, T.:* Two dimensional Berezin-Li-Yau inequalities with a correction term
- 2007-006 *Weidl, T.:* Improved Berezin-Li-Yau inequalities with a remainder term
- 2007-005 *Frank, R.L.; Loss, M.; Weidl, T.:* Polyá's conjecture in the presence of a constant magnetic field
- 2007-004 *Ekholm, T.; Frank, R.L.; Kovarik, H.:* Eigenvalue estimates for Schrödinger operators on metric trees
- 2007-003 *Lesky, P.H.; Racke, R.:* Elastic and electro-magnetic waves in infinite waveguides
- 2007-002 *Teufel, E.:* Spherical transforms and Radon transforms in Moebius geometry
- 2007-001 *Meister, A.:* Deconvolution from Fourier-oscillating error densities under decay and smoothness restrictions